

REPRESENTABILITY OF CONES OF MONOTONE FUNCTIONS IN WEIGHTED LEBESGUE SPACES AND EXTRAPOLATION OF OPERATORS ON THESE CONES

E. I. BEREZHNOÏ AND L. MALIGRANDA

ABSTRACT. It is shown that a sublinear operator is bounded on the cone of monotone functions if and only if a certain new operator related to the one mentioned above is bounded on a certain ideal space defined constructively. This construction is used to provide new extrapolation theorems for operators on the cone in weighted Lebesgue spaces.

§1. INTRODUCTION

The role of sharp estimates for classical operators in harmonic analysis and related fields is well known. In the recent time, in connection with new analytic problems, it has become fashionable to estimate such operators on certain cones in a given space rather than on the entire space (see, e.g., [1, 4, 17, 20, 27, 28, 36, 38, 40, 41]). Next, for operators with positive kernels the Schur extrapolation theorem is also well known (see, e.g., [25]), saying that an integral operator $Tx(t) = \int k(t, s)x(s) ds$, $k(t, s) \geq 0$ is bounded on L^p if and only if there exists a positive function $u(t)$ finite a.e. and such that the operator is bounded in the couples $T: L_u^\infty \rightarrow L_u^\infty$ and $T: L_v^1 \rightarrow L_v^1$, where $v = u^{1/p-1}$. Since various problems of analysis have resulted in a gradually increasing interest to extrapolation theorems, see [5, 7–9], it seems to be natural to pass from spaces to cones in the extrapolation theory for L^p .

The present work was planned as early as in the beginning of the 2000s. A short summary of the main results was given in [10]. The central result of the paper consists of the verification of the fact that, basically, the cone $K(\downarrow) \cap L_v^p$ in the Lebesgue space L_v^p is generated by the linear operator $Qx(t) = \int_t^\infty x(s) ds$ of integration. For the operators T in the class $\text{Sub}(\alpha, \beta, \gamma, \downarrow)$ (which is described below and contains all subadditive operators), this makes it possible to prove the equivalence $T: K(\downarrow) \cap L_v^p \rightarrow X \Leftrightarrow TQ: L_{\bar{v}}^p \rightarrow X$. Here the weight \bar{v} is defined constructively in terms of the weight function v (see Theorem 1). This approach distinguishes our paper from the well-known paper [16], which is devoted to estimates of classical operators in the couple $(K(\downarrow) \cap L_v^p, L_w^p)$ implied by certain estimates in couples of weighted Lebesgue spaces.

Our approach allows us to apply the entire technique of sharp estimates on weighted Lebesgue spaces to the derivation of sharp estimates of operators on cones. In particular, these constructions have led us to a new extrapolation theorem for operators in the class $\text{Sub}(\alpha, \beta, \gamma, \downarrow)$ that are defined on cones included in $K(\downarrow) \cap L_v^p$ in a weighted Lebesgue space. This extrapolation theorem is new even for the Hardy classical operator.

2010 *Mathematics Subject Classification.* Primary 46E30, 46B20, 46B42.

Key words and phrases. Weighted Lebesgue space, cone of monotone functions, extrapolation of operators.

Supported by RFBR (grant no. 14-01-00417).

§2. PRELIMINARIES

Let $S(\mu) = S(R_+, \Sigma, \mu)$ ($R_+ = (0, +\infty)$) denote the space of measurable functions $x: R_+ \rightarrow R$, let $\chi(D)$ stand for the characteristic function of a set D , and let $\|x\|_X$ be the norm of an element x in X . Recall that a Banach space $X = (X, \|\cdot\|_X)$ of measurable functions is called an ideal space (see [24, 26]) if $x \in X$ and $\|x\|_X \leq \|y\|_X$ whenever x is measurable and $|x(t)| \leq |y(t)|$ a.e. on R_+ for some $y \in X$. As usual, the symbol L^p ($1 \leq p \leq \infty$) denotes the Lebesgue space, and the exponent conjugate to $p \in [1, \infty]$ is denoted by p' : $\frac{1}{p} + \frac{1}{p'} = 1$.

Let $v: R_+ \rightarrow R_+$ be a positive function (a weight). If X is an ideal space, we denote by X_v the new ideal space whose norm is defined as follows: $\|x\|_{X_v} = \|vx\|_X$. In particular, the norm in L_v^p ($1 \leq p < \infty$) has the form

$$\|x\|_{L_v^p} = \left(\int_0^\infty |x(t)v(t)|^p dt \right)^{1/p},$$

which differs somewhat from the variant usually adopted (the latter presumes the incorporation of the weight in the measure).

For every ideal space X , the dual space X' is well defined. It consists of bounded linear functionals on X representable by integrals; the norm of every such functional is defined by $\|y\|_{X'} = \sup \left\{ \int_R y(t)x(t) dt : \|x\|_X \leq 1 \right\}$. If v is a weight and X is an ideal space, it can easily be verified that

$$(1) \quad (X_v)' = (X')_{1/v}.$$

Let X be an ideal space in $S(\mu)$, and K a cone in $S(\mu)$. The symbol $K \cap X$ denotes the intersection of K with X_+ .

Definition 1. We denote by $K(\downarrow)$ the cone in $S(\mu)$ consisting of monotone nonincreasing functions $x: R_+ \rightarrow R_+$, i.e., $x(t+h) \leq x(t)$ for $h \geq 0$. Similarly, $K(\uparrow)$ denotes the cone of monotone nondecreasing functions in $S(\mu)$.

Now, we describe the classes of operators to be treated in the paper.

We denote by $\text{Sub}(+)$ the class of sublinear operators T defined on an ideal space X or on $S(\mu)$ and taking values in $S(\mu)$. For $T \in \text{Sub}(+)$, the adjoint operator may fail to exist, but its role can be played by the associated operator $T' \in \text{Sub}(+)$ defined as follows.

For $T \in \text{Sub}(+)$, an operator $T' \in \text{Sub}(+)$ is said to be *associated* with T in the scale L^p if for all $1 \leq p \leq \infty$ and all weight functions u , the operator $T: L_u^p \rightarrow L_u^p$ is bounded if and only if $T': L_{1/u}^{p'} \rightarrow L_{1/u}^{p'}$ is also bounded and, moreover,

$$C^{-1} \|T\|_{L_u^p \rightarrow L_u^p} \leq \|T'\|_{L_{1/u}^{p'} \rightarrow L_{1/u}^{p'}} \leq C \|T\|_{L_u^p \rightarrow L_u^p}$$

with some constant $C > 0$ independent of p and u .

An associated operator is not uniquely determined. If T is linear, we may take the adjoint T^* for the role of an associated operator T' . Next, for a linear operator T , the operator $x \mapsto |Tx|$ is sublinear and possesses no adjoint, but the operator $T'x = |T^*x|$ is associated with it. If $T \in \text{Sub}(+)$ and a linear operator T_1 is given, then the role of operators associated with the compositions TT_1 and T_1T can be played by $(T_1)^*T'$ and $T'(T_1)^*$. Thus, the set $T, T' \in \text{Sub}(+)$ is a two-sided ideal with respect to composition with bounded linear operators.

Now, we extend the class $\text{Sub}(+)$.

Definition 2. We say that an operator $T: X \cap K(\downarrow) \rightarrow Y$ belongs to the class $\text{Sub}(\alpha, \beta, \gamma, \downarrow)$, ($\alpha \geq 1, \beta > 0, \gamma > 0$) if

a) for every $x, y \in X \cap K(\downarrow)$ we have

$$\|T(y+x) \mid Y\| \leq \alpha(\|Ty \mid Y\| + \|Tx \mid Y\|);$$

b) for every $x \in X \cap K(\downarrow)$ and every $\lambda \in R$ we have

$$\|T(\lambda x) \mid Y\| = |\lambda| \|Tx \mid Y\|;$$

c) for every $x \in X \cap K(\downarrow)$ we have

$$\inf \{ \|Ty \mid Y\| : y(t) \geq \beta x(t) : y \in X \cap K(\downarrow) \} \geq \gamma \|Tx \mid Y\|.$$

It is straightforward from the definition that every operator belonging to $\text{Sub}(+)$ belongs also to $\text{Sub}(\downarrow)$. To see this, it suffices to put $\alpha = 1$, choose a positive number β arbitrarily, and define γ by $\gamma = \max\{1, \frac{1}{\beta}\}$.

For $T \in \text{Sub}(\alpha, \beta, \gamma, \downarrow)$, we can define an operator T' associated with it in the scale L^p by analogy with the case of $T \in \text{Sub}(+)$.

The proof of the following lemma is an easy consequence of the definitions.

Lemma 1. (a) Suppose that $T \in \text{Sub}(\alpha, \beta, \gamma, \downarrow)$ and $\delta \in (0, \infty)$. Then

(a) if $\delta > 1$, then $T \in \text{Sub}(\alpha, \delta\beta, \gamma, \downarrow)$;

(b) if $\delta < 1$, then $T \in \text{Sub}(\alpha, \delta\beta, \frac{1}{\delta}\gamma, \downarrow)$.

Since precise values of the constants are irrelevant in the present paper, we introduce the notation $\text{Sub}(\downarrow)$ for the following class of operators:

$$\text{Sub}(\downarrow) = \bigcup_{\beta, \gamma > 0} \left(\bigcup_{\alpha \geq 1} \text{Sub}(\alpha, \beta, \gamma, \downarrow) \right).$$

We present an example showing that $\text{Sub}(\downarrow)$ is much wider than $\text{Sub}(+)$.

Fix a monotone increasing sequence $\{k_i\}_1^\infty$ of positive integers; let $k_1 > 4$. We introduce a function $w: [0, \infty) \rightarrow R_+$. On each interval $[i, i+1)$ it is given by

$$w(t) = \begin{cases} 1, & \text{for } t \in [i-1, i - \frac{1}{k_i}), \\ -\frac{k_i}{4}, & \text{for } t \in [i - \frac{1}{k_i}, i), \end{cases} \quad i = 1, 2, \dots$$

Now, we define a functional f by the formula

$$f(x) = \int_0^\infty w(s)x(s) ds.$$

Then for every $x \in K(\downarrow)$ we have

$$\begin{aligned} \int_{i-1}^i w(s)x(s) ds &= \int_{i-1}^{i-\frac{1}{k_i}} w(s)x(s) ds + \int_{i-\frac{1}{k_i}}^i w(s)x(s) ds \\ &\geq \int_{i-1}^{i-\frac{1}{k_i}} x(s) ds - k_i \frac{1}{4k_i} x\left(i - \frac{1}{k_i}\right) \\ &\geq \int_{i-1}^{i-\frac{1}{2}} x(s) ds \geq \frac{1}{2} \int_{i-1}^i x(s) ds. \end{aligned}$$

This inequality shows that

$$\begin{aligned} \int_0^\infty x(s) ds &\geq \int_0^\infty w(s)x(s) ds = f(x) = \sum_{i=1}^\infty \int_{i-1}^i w(s)x(s) ds \\ &\geq \frac{1}{2} \sum_{i=1}^\infty \int_{i-1}^i x(s) ds = \frac{1}{2} \int_0^\infty x(s) ds \end{aligned}$$

for every $x \in K(\downarrow)$. But now, if $y \in K(\downarrow)$ and $y(t) \geq \beta x(t)$ for a.e. t , then, applying the last inequality, we obtain

$$f(x) \leq \int_0^\infty x(s) ds \leq \frac{1}{\beta} \int_0^\infty y(s) ds \leq \frac{2}{\beta} f(y).$$

Thus, the functional f constructed above belongs to $\text{Sub}(1, \beta, \frac{2}{\beta}, \downarrow)$. On the other hand, since $\lim_{i \rightarrow \infty} k_i = \infty$, we see that there is no nonnegative function w_0 having the property that the functionals

$$f(x) = \int_0^\infty w(s)x(s) ds, \quad f_0(x) = \int_0^\infty w_0(s)x(s) ds$$

are equivalent on the cone of nonnegative functions. Thus, the functional f is not equivalent to any functional f_0 belonging to $\text{Sub}(+)$.

Below we shall often use the classical integral operators given by the following formulas on their natural domains:

$$Px(t) = \int_0^t x(s) ds, \quad Qx(t) = \int_t^\infty x(s) ds.$$

The next result about the boundedness of these operators in weighted $(L^p - L^q)$ -spaces is well known (see [28, 35] and [27]).

Lemma 2. (a) Let $1 \leq p \leq q \leq \infty$. Then the operator $P: L_v^p \rightarrow L_w^q$ is bounded if and only if

$$(2) \quad \sup_{t>0} \left\| \frac{1}{v} \chi_{[0,t]} \mid L^{p'} \right\| \|w \chi_{[t,\infty)} \mid L^q\| < \infty.$$

The operator $Q: L_v^p \rightarrow L_w^q$ is bounded if and only if

$$(3) \quad \sup_{t>0} \left\| \frac{1}{v} \chi_{[t,\infty)} \mid L^{p'} \right\| \|w \chi_{[0,t]} \mid L^q\| < \infty.$$

(b) Let $1 < q < p < \infty$, $\frac{1}{r} = \frac{1}{q} - \frac{1}{p}$. Then the operator $P: L_v^p \rightarrow L_w^q$ is bounded if and only if

$$(4) \quad \left(\int_0^\infty \left(\left\| \frac{1}{v} \chi_{[0,t]} \mid L^{p'} \right\|^{p'/q'} \|w \chi_{[t,\infty)} \mid L^q\| \right)^r v(t)^{-p'} dt \right)^{1/r} < \infty.$$

The operator $Q: L_v^p \rightarrow L_w^q$ is bounded if and only if

$$(5) \quad \left(\int_0^\infty \left(\left\| \frac{1}{v} \chi_{[t,\infty)} \mid L^{p'} \right\|^{p'/q'} \|w \chi_{[0,t]} \mid L^q\| \right)^r v(t)^{-p'} dt \right)^{1/r} < \infty.$$

(c) If $1 = q < p < \infty$, we have $r = p'$. Therefore, formula (4) should be understood in the following way:

$$(6) \quad \left(\int_0^\infty \|w \chi_{[t,\infty)} \mid L^1\|^{p'} v(t)^{-p'} dt \right)^{1/p'} < \infty.$$

Similarly, a limit passage in (5) yields

$$(7) \quad \left(\int_0^\infty \|w \chi_{[0,t]} \mid L^1\|^{p'} v(t)^{-p'} dt \right)^{1/p'} < \infty.$$

(d) If $1 \leq q < p = \infty$, formula (4) becomes

$$(8) \quad \left(\int_0^\infty \left(w(t) \left\| \frac{1}{v} \chi_{[0,t]} \mid L^1 \right\| \right)^q dt \right)^{1/q} < \infty,$$

and inequality (5) transforms to

$$(9) \quad \left(\int_0^\infty \left(w(t) \left\| \frac{1}{v} \chi_{[t,\infty)} \mid L^1 \right\| \right)^q dt \right)^{1/q} < \infty.$$

§3. REPRESENTATION OF CONES IN WEIGHTED L^p -SPACES

We begin with the statements of two main results (Theorems 1 and 2), which constitute a principal tool for the study of operators belonging to $\text{Sub}(\downarrow)$ on cones. We begin with the case where $1 \leq p < \infty$.

Theorem 1. Fix a number $1 \leq p < \infty$ and a weight function v such that

$$(10) \quad \int_0^t v(s)^p ds < \infty$$

for every $t \in \mathbb{R}_+$ and

$$(11) \quad \int_0^\infty v(s)^p ds = \infty.$$

We introduce a new weight function \bar{v} , putting

$$(12) \quad \|v \chi_{[0,t]} \mid L^p\| \left\| \frac{1}{v} \chi_{[t,\infty)} \mid L^{p'} \right\| \equiv 1.$$

Then

(a) $Q((L_{\bar{v}}^p)_+) \subset K(\downarrow) \cap L_v^p$ and, moreover, for every $x \in (L_{\bar{v}}^p)_+$ we have

$$\|Qx \mid L_v^p\| \leq \|x \mid L_{\bar{v}}^p\|;$$

(b) for every $x \in K(\downarrow) \cap (L_v^p)_+$ and every $\varepsilon > 0$ there exists $x_\varepsilon \in (L_{\bar{v}}^p)_+$ such that

$$(13) \quad \|x_\varepsilon \mid L_{\bar{v}}^p\| \leq 16(1 + \varepsilon) \|x \mid L_v^p\|$$

and

$$(14) \quad Q(x_\varepsilon)(t) \geq \frac{1}{8} x(t)$$

for a.e. $t > 0$.

The proof of Theorem 1 will be given in the last section of the paper. Here we comment on its assumptions and show some applications.

The assumption (10) says that for every $t > 0$ the characteristic function $\chi_{[0,t]}$ satisfies the condition $\chi_{[0,t]} \in K(\downarrow) \cap L_v^p$, i.e., the cone $K(\downarrow) \cap L_v^p$ is nondegenerate. The assumption (11) says that every $x \in K(\downarrow) \cap L_v^p$ satisfies $\lim_{t \rightarrow \infty} x(t) = 0$. It should be noted that if $\int_0^\infty v(s)^p ds < \infty$, then for $p \in [1, \infty)$ relation (12) must fail as $t \rightarrow +\infty$ or as $t \rightarrow \infty$.

Relation (12) for \bar{v} can be expanded as follows:

$$(15) \quad \bar{v}(t) = \frac{(p-1)^{1/p'}}{v(t)^{p-1}} \int_0^t v(s)^p ds, \quad p > 1, \quad \bar{v}(t) = \int_0^t v(s) ds \quad \text{for } p = 1.$$

Now we present a series of corollaries to Theorem 1.

Corollary 1. Let $1 \leq p \leq q \leq \infty$ ($p \neq \infty$), and let the weight function v satisfy conditions (10), (11). Then the embedding $K(\downarrow) \cap (L_v^p)_+ \subset (L_w^q)_+$ (equivalently, the inequality

$$(16) \quad \|x \mid L_w^q\| \leq C_1 \|x \mid L_v^p\|$$

for every $x \in K(\downarrow) \cap (L_v^p)_+$ occurs if and only if

$$(17) \quad \sup_{t>0} \frac{\|w\chi_{[0,t]} \mid L^q\|}{\|v\chi_{[0,t]} \mid L^p\|} = C_2 < \infty.$$

Proof. The necessity of condition (17) follows because $\chi_{[0,t]} \in K(\downarrow) \cap (L_v^p)_+$. To prove sufficiency, we use Theorem 1: condition (16) is fulfilled if and only if

$$(18) \quad \|Qx \mid L_w^q\| \leq C_3 \|x \mid L_v^p\|$$

for every $x \in (L_v^p)_+$, where the weight function \bar{v} is defined in (12). Applying (17) and (12), we obtain

$$\infty > C_2 = \sup_{t>0} \frac{\|w\chi_{[0,t]} \mid L^q\|}{\|v\chi_{[0,t]} \mid L^p\|} = \sup_{t>0} \|w\chi_{[0,t]} \mid L^q\| \left\| \frac{1}{\bar{v}} \chi_{[t,\infty)} \mid L^{p'} \right\|.$$

The last inequality and Lemma 2 yield (18). \square

Proofs of Corollary 1 based on different ideas can be found in the papers by Sawyer (see [36, Remark (i), p. 148]), Stepanov (see [41, Proposition 1(a)]), Carro and Soria (see [14, Corollary 2.7]), and Heinig and Maligranda (see [20, Proposition 2.5(a)]). The structure of the cone $K(\downarrow) \cap L_v^p$ considered in the Lorentz quasi-Banach space Λ_{p,v^p} was also treated in [22].

Corollary 2. Let $1 \leq q < p < \infty$, $\frac{1}{r} = \frac{1}{q} - \frac{1}{p}$, and let a weight function v satisfy conditions (10) and (11). Then the embedding

$$(19) \quad K(\downarrow) \cap (L_v^p)_+ \subset (L_w^q)_+$$

or, equivalently, the relation

$$\|x \mid L_w^q\| \leq C_1 \|x \mid L_v^p\| \text{ for every } x \in K(\downarrow) \cap (L_v^p)_+,$$

occurs if and only if

$$(20) \quad C_4 := \left(\int_0^\infty (\|w\chi_{[0,t]} \mid L^q\|)^r \|v\chi_{[0,t]} \mid L^p\|^{-pr/q} v^p(t) dt \right)^{1/r} < \infty.$$

Proof. Condition (19) is fulfilled if and only if the identity operator I maps boundedly the cone $K(\downarrow) \cap L_v^p$ to the cone $(L_w^q)_+$. We show that the latter is equivalent to the following relation for Q :

$$(21) \quad \|Q \mid L_v^p \rightarrow L_w^q\| < \infty.$$

To see that (21) suffices, we apply (13) and (14):

$$\begin{aligned} \sup \{ \|x \mid L_w^q\| : x \in K(\downarrow) \cap L_v^p \text{ \& } \|x \mid L_v^p\| \leq 1 \} \\ \leq 8 \sup \{ \|x \mid L_w^q\| : x \leq Qx_\varepsilon \text{ \& } \|x_\varepsilon \mid L_v^p\| \leq 16(1+\varepsilon) \}. \end{aligned}$$

The necessity of (21) follows from (12)–(14):

$$\|Q \mid L_v^p \rightarrow L_w^q\| = \|I(Q) \mid L_v^p \rightarrow L_w^q\| \leq \|Q \mid L_v^p \rightarrow L_v^p\| \|I \mid K(\downarrow) \cap L_v^p \rightarrow L_w^q\|.$$

By (5), condition (21) is equivalent to the inequality

$$\left(\int_0^\infty \left(\left\| \frac{1}{\bar{v}} \chi_{[t,\infty)} \mid L^{p'} \right\|^{p'/q'} \|w\chi_{[0,t]} \mid L^q\| \right)^r \bar{v}(t)^{-p'} dt \right)^{1/r} < \infty$$

or

$$(22) \quad \left(\int_0^\infty (\|w\chi_{[0,t]} \mid L^q\| \|v\chi_{[0,t]} \mid L^p\|^{-p'/q'})^r \left(-\frac{d}{dt} \|v\chi_{[0,t]} \mid L^p\|^{-p'} \right) dt \right)^{1/r} < \infty.$$

Since

$$\begin{aligned} -\frac{d}{dt}\|v\chi_{[0,t]}|L^p\|^{-p'} &= -\frac{d}{dt}\left(\int_0^t v^p(s)ds\right)^{(1-p')} \\ &= (p'-1)\left(\int_0^t v^p(s)ds\right)^{-p'}v^p(t) = (p'-1)\|v\chi_{[0,t]}|L^p\|^{-pp'}v^p(t), \end{aligned}$$

we see that (22) is a consequence of the relations

$$\begin{aligned} &\left(\int_0^\infty (\|w\chi_{[0,t]}|L^q\|\|v\chi_{[0,t]}|L^p\|^{-p'/q'})^r\left(-\frac{d}{dt}\|v\chi_{[0,t]}|L^p\|^{-p'}\right)dt\right)^{1/r} \\ &= \left((p'-1)\int_0^\infty \|w\chi_{[0,t]}|L^q\|^r\|v\chi_{[0,t]}|L^p\|^{-(p'p+\frac{p'r}{q'})}v^p(t)dt\right)^{1/r} \\ &= \left((p'-1)\int_0^\infty \|w\chi_{[0,t]}|L^q\|^r\|v\chi_{[0,t]}|L^p\|^{-pr/q}v^p(t)dt\right)^{1/r} \\ &= (p'-1)^{1/r}\cdot C_4. \end{aligned} \quad \square$$

Other proofs of Corollary 2 can be found in the papers by Sawyer (see [36, Remark (i), p. 148]) and Stepanov (see [41, Proposition 1(b)]).

Corollary 3. *Let $1 \leq q < p < \infty$, $\frac{1}{r} = \frac{1}{q} - \frac{1}{p}$, and suppose that a weight function v satisfies conditions (10), (11) for p and a weight function w satisfies conditions (10), (11) for q . Then*

$$K(\downarrow) \cap (L_v^p)_+ \neq K(\downarrow) \cap (L_w^q)_+,$$

that is, for every v, w with (10) and (11) these two cones do not coincide.

Proof. Suppose the contrary, i.e., let

$$K(\downarrow) \cap (L_v^p)_+ = K(\downarrow) \cap (L_w^q)_+.$$

Then, by Corollary 2, the embedding $K(\downarrow) \cap (L_v^p)_+ \subset (L_w^q)_+$ is equivalent to (21) or (22):

$$(23) \quad \left(\int_0^\infty (\|w\chi_{[0,t]}|L^q\|\|v\chi_{[0,t]}|L^p\|^{-p'/q'})^r\left(-\frac{d}{dt}\|v\chi_{[0,t]}|L^p\|^{-p'}\right)dt\right)^{1/r} < \infty.$$

Next, by Corollary 1, the embedding $K(\downarrow) \cap (L_w^q)_+ \subset (L_v^p)_+$ is equivalent to (17):

$$(24) \quad \sup_{t>0} \frac{\|v\chi_{[0,t]}|L^p\|}{\|w\chi_{[0,t]}|L^q\|} = C_2 < \infty.$$

Then, using (23) and (24), we obtain

$$\begin{aligned} &\infty > \left(\int_0^\infty (\|w\chi_{[0,t]}|L^q\|\|v\chi_{[0,t]}|L^p\|^{-p'/q'})^r\left(-\frac{d}{dt}\|v\chi_{[0,t]}|L^p\|^{-p'}\right)dt\right)^{1/r} \\ &\geq \frac{1}{C_2}\left(\int_0^\infty (\|v\chi_{[0,t]}|L^p\|\|v\chi_{[0,t]}|L^p\|^{-p'/q'})^r\left(-\frac{d}{dt}\|v\chi_{[0,t]}|L^p\|^{-p'}\right)dt\right)^{1/r}. \end{aligned}$$

Recalling that $(1 - \frac{p'}{q'})r = \frac{p}{p-1} = p'$ and substituting $\tau = \|v\chi_{[0,t]}|L^p\|^{-p'}$, which is possible because

$$\lim_{t \rightarrow 0} \|v\chi_{[0,t]}|L^p\|^{p'} = 0, \quad \lim_{t \rightarrow \infty} \|v\chi_{[0,t]}|L^p\|^{p'} = \infty,$$

we arrive at

$$\infty > \frac{1}{C_2}\left(\int_\infty^0 \frac{1}{\tau}(-d\tau)\right)^{1/r} = \frac{1}{C_2}\left(\int_0^\infty \frac{1}{\tau}d\tau\right)^{1/r} = \infty.$$

This contradiction shows that the cones do not coincide. \square

Now, we present a result about norm estimates for operators in $\text{Sub}(\downarrow)$.

Theorem 2. *Fix a number $p \in [1, \infty)$, take a weight function v satisfying (10), (11), and construct the function \bar{v} as in Theorem 1. Let X be an ideal space.*

An operator $T \in \text{Sub}(\downarrow)$ acts boundedly from $K(\downarrow) \cap L_v^p$ to X , i.e.,

$$(25) \quad \|Tx \mid X\| \leq C_5 \|x \mid L_v^p\|$$

for every $x \in K(\downarrow) \cap (L_v^p)_+$, if and only if the composition operator TQ acts boundedly from $L_{\bar{v}}^p$ to X , i.e.,

$$(26) \quad \|TQx \mid X\| \leq C_6 \|x \mid L_{\bar{v}}^p\|$$

for every $x \in (L_{\bar{v}}^p)_+$.

Proof. First, we show that (26) \implies (25). By Theorem 1, for every $x \in K(\downarrow) \cap (L_v^p)_+$ there exists $x_\varepsilon \in (L_v^p)_+$ such that

$$\|x_\varepsilon \mid L_{\bar{v}}^p\| \leq 16(1 + \varepsilon) \|x \mid L_v^p\| \text{ and } Q(x_\varepsilon)(t) \geq \frac{1}{8}x(t) \text{ for all } t > 0.$$

The definition of the set $\text{Sub}(\downarrow)$ and Lemma 1 imply the existence of constants $\alpha \geq 1$ and $\gamma > 0$ with $T \in \text{Sub}(\alpha, \frac{1}{8}, \gamma; \downarrow)$. Therefore,

$$\|Tx \mid X\| \leq \gamma \|TQ(x_\varepsilon) \mid X\| \leq \gamma C_6 \|x_\varepsilon \mid L_{\bar{v}}^p\| \leq 16(1 + \varepsilon) \gamma C_6 \|x \mid L_v^p\|,$$

which proves the implication (26) \implies (25).

Now, we verify the reverse implication: (25) \implies (26). The mapping Q takes any nonnegative function to a monotone nonincreasing function, i.e., $Qx \in K(\downarrow) \cap (L_v^p)_+$ for every $x \in (L_v^p)_+$. Next, by the definition (12) of \bar{v} , the operator Q is bounded when treated as an operator $Q: L_v^p \rightarrow L_{\bar{v}}^p$. Therefore, we have

$$\|TQx \mid X\| \leq C_5 \|Qx \mid L_v^p\| \leq C_5 \|Q \mid L_v^p \rightarrow L_{\bar{v}}^p\| \|x \mid L_v^p\|. \quad \square$$

Using the techniques of estimating operators $L: L_w^p \rightarrow X$ (see, e.g., [3, 4, 6, 27, 28]), on the basis of Theorem 2 it is possible to deduce various estimates for operators on the cone of monotone functions in Lebesgue spaces. We illustrate this by several classical examples.

First, with the help of a new approach, we shall prove the theorem of Sawyer (see [36]), which, in combination with a result by Ariño and Muckenhoupt (see [1]), resolved an important problem of harmonic analysis, namely, the boundedness problem for the Hardy operator on weighted Lorentz spaces. Furthermore, that theorem gave rise to a wide range of new problems, which remain fashionable still.

Theorem 3. *Let p, v , and \bar{v} be the same as in Theorem 1. Consider a measurable function $g: R_+ \rightarrow R_+$. Then*

$$(27) \quad \frac{1}{C_8} \left\| \int_0^t g(s) ds \mid L_{\frac{1}{\bar{v}}}^{p'} \right\| \leq \sup \left\{ \int_0^\infty y(t)g(t) dt : y \in K(\downarrow) \cap L_v^p, \|y \mid L_v^p\| \leq 1 \right\} \\ \leq C_8 \left\| \int_0^t g(s) ds \mid L_{\frac{1}{\bar{v}}}^{p'} \right\|,$$

where the constant $C_8 > 0$ does not depend on g .

Proof. We define a functional $F: K(\downarrow) \cap L_v^p \rightarrow R$ by $Fy(t) = \int_0^\infty y(t)g(t) dt$. Since this functional is quasilinear and nonnegative, we may apply Theorem 2 to deduce the existence of a constant $c > 0$ such that

$$\frac{1}{c} \|FQ \mid L_{\bar{v}}^{p'} \rightarrow R\| \leq \|F \mid K(\downarrow) \cap L_v^p \rightarrow R\| \leq c \|FQ \mid L_{\bar{v}}^{p'} \rightarrow R\|.$$

Integrating by parts, we arrive at

$$(28) \quad \int_0^\infty g(t)Qy(t) dt = \int_t^\infty y(s) ds \int_0^t g(s) ds \Big|_0^\infty + \int_0^\infty \left(\int_0^t g(s) ds \right) y(t) dt.$$

Suppose first that

$$(29) \quad \lim_{t \rightarrow 0} \frac{1}{\|\chi_{[0,t]}v\|_{L^p}} \cdot \int_0^t g(s) ds = 0, \quad \lim_{t \rightarrow \infty} \frac{1}{\|\chi_{[0,t]}v\|_{L^p}} \cdot \int_0^t g(s) ds = 0.$$

Then, whenever $y \in K(\downarrow) \cap L^p_v$ satisfies $\|y\|_{K(\downarrow) \cap L^p_v} \leq 1$, by (1) we can estimate the integrated term as follows:

$$\left| \int_t^\infty y(s) ds \int_0^t g(s) ds \right| \leq \left\| \chi_{[t,\infty)} \frac{1}{v} \right\|_{L^{p'}} \cdot \int_0^t g(s) ds = \frac{1}{\|\chi_{[0,t]}v\|_{L^p}} \cdot \int_0^t g(s) ds.$$

Now, (27) is a consequence of (28), the last inequality, (29), and the definition of the dual norm.

The assumption (29) can be lifted in the following way.

For a nonnegative function g and arbitrary $n \in \mathbb{N}$, put $g_n(t) \equiv \chi_{(n^{-1},n)}(t)g(t)$. Then g_n satisfies (29), whence we obtain (27). We can easily pass to the limit here with the help of the B. Levy classical theorem (see, e.g., [24]). \square

Note that, tracing the behavior of the constant in Theorem 3, it is possible to estimate the constant in (27).

Now we consider one of the most important operators in analysis, namely, the Hardy operator. On its natural domain, it is defined by the formula

$$Hx(t) = \frac{1}{t} \int_0^t x(s) ds.$$

Theorem 4. Suppose that $1 \leq p < \infty$, $1 \leq q \leq \infty$, and functions v and \bar{v} satisfy the assumptions of Theorem 1. For the Hardy operator to be bounded in the sense that

$$(30) \quad H: K(\downarrow) \cap L^p_v \rightarrow L^q_w,$$

it is necessary and sufficient that the following conditions be fulfilled:

a) if $1 \leq p \leq q \leq \infty$, $\frac{1}{p} + \frac{1}{p'} = 1$, then

$$(31) \quad \sup_{t>0} \frac{\|\chi_{(0,t)}w(s)\|_{L^q}}{\|\chi_{(0,t)}v(s)\|_{L^p}} < \infty; \quad \sup_{t>0} \frac{1}{t} \left\| \chi_{(0,t)} \frac{s}{\bar{v}(s)} \right\|_{L^{p'}} \cdot \|\chi_{(t,\infty)}w(s)\|_{L^q} < \infty;$$

b) if $1 \leq q < p < \infty$, $\frac{1}{r} = \frac{1}{q} - \frac{1}{p}$, then

$$(32) \quad \left(\int_0^\infty \left(\frac{\|\chi_{(0,t)}w(s)\|_{L^q}}{\|\chi_{(0,t)}v(s)\|_{L^p}} \right)^r \left(-\frac{d}{dt} \left(\frac{1}{\|\chi_{(t,\infty)}v(s)\|_{L^p}} \right)^{p'} \right) dt \right)^{1/r} < \infty;$$

$$(33) \quad \left(\int_0^\infty \left(\left\| \chi_{(0,t)} \frac{s}{\bar{v}(s)} \right\|_{L^{p'}} \cdot \|\chi_{(t,\infty)}w(s)\|_{L^q} \right)^r \left(-\frac{d}{dt} \left(\frac{1}{\|\chi_{(t,\infty)}v(s)\|_{L^p}} \right)^{p'} \right) dt \right)^{1/r} < \infty.$$

Proof. By Theorem 2, relation (30) is fulfilled if and only if the operator HQ acts boundedly in the couple

$$HQ: L^p_v \rightarrow L^q_w.$$

Since HQ is positive, it suffices to verify its boundedness on nonnegative functions. Let $x(t) \geq 0$ a.e. Using the Fubini theorem (see, e.g., [24]) for nonnegative functions, we

obtain

$$\begin{aligned} HQx(t) &= \frac{1}{t} \int_0^t \left(\int_s^\infty x(\tau) d\tau \right) ds \\ &= \frac{1}{t} \left(\int_0^t \left[\int_s^t x(\tau) d\tau + \int_t^\infty x(\tau) d\tau \right] ds \right) \\ &= \frac{1}{t} \left(\int_0^t \left[\int_0^\tau ds \right] x(\tau) d\tau + t \int_t^\infty x(\tau) d\tau \right) \\ &= \frac{1}{t} \int_0^t \tau x(\tau) d\tau + \int_t^\infty x(\tau) d\tau. \end{aligned}$$

The proof can be finished by application of Lemma 2 to each summand in the last identity. \square

Now we pass to the cone $K(\downarrow) \cap L_v^\infty$. Note that the situation will differ much from the case of $p < \infty$ considered above.

Our goal is to present certain analogs of the statements formulated above for the cone $K(\downarrow) \cap L_v^\infty$. Despite the relative ease of proofs, the central results of this subsections are Theorems 5 and 6.

Theorem 5. *Fixing $p = \infty$ and a weight function $v: [0, \infty) \rightarrow R_+$, we define a new function \tilde{v} by the formula*

$$(34) \quad \tilde{v}(t) = \operatorname{ess\,sup}_{0 < \tau < t} v(\tau).$$

Then \tilde{v} is monotone nondecreasing, and the cones $K(\downarrow) \cap L_v^\infty$ and $K(\downarrow) \cap L_{\tilde{v}}^\infty$ coincide. Moreover, for every $x \in K(\downarrow) \cap L_v^\infty$ we have

$$(35) \quad \|x\|_{L_v^\infty} = \|x\|_{L_{\tilde{v}}^\infty}.$$

Proof. The definition (34) readily implies that the function \tilde{v} is monotonic.

We verify (35). Directly from the definition (34), it follows that for a.e. $t \in [0, \infty)$ we have $\tilde{v}(t) \geq v(t)$. Consequently, for every $x \in K(\downarrow) \cap L_v^\infty$ we obtain the norm inequality

$$(36) \quad \|x\|_{L_{\tilde{v}}^\infty} \geq \|x\|_{L_v^\infty}.$$

Now, let $x \in K(\downarrow)$. Then

$$\tilde{v}(t)x(t) \leq \operatorname{ess\,sup}_{0 < s \leq t} v(s)x(s) \leq \operatorname{ess\,sup}_{s > 0} v(s)x(s) = \|x\|_{L_v^\infty}$$

for a.e. $t \in R_+$. Thus,

$$(37) \quad \|x\|_{L_v^\infty} \leq \|x\|_{L_{\tilde{v}}^\infty}.$$

By (36)–(37), identity (35) follows. \square

Corollary 4. *Let $p = \infty$, $q \in [1, \infty]$. Then the embedding $K(\downarrow) \cap (L_v^\infty)_+ \subset (L_w^q)_+$ or, equivalently, the inequality*

$$(38) \quad \|x\|_{L_w^q} \leq C_1 \|x\|_{L_v^\infty}$$

for every $x \in K(\downarrow) \cap (L_v^\infty)_+$, occur if and only if

$$(39) \quad \left\| \frac{1}{\tilde{v}} w \right\|_{L^q} = C_1 < \infty.$$

Proof. Theorem 5 shows that the cones $K(\downarrow) \cap L_v^\infty$ and $K(\downarrow) \cap L_{\tilde{v}}^\infty$ coincide. The function $\frac{1}{\tilde{v}}$, which belongs to the intersection of $K(\downarrow)$ and the unit ball of $L_{\tilde{v}}^\infty$, is a pointwise majorant for all functions in $K(\downarrow) \cap L_{\tilde{v}}^\infty$ with unit norm. Thus, conditions (38) and (39) are equivalent. \square

Corollary 5. *Let $p = \infty$.*

If $q \neq \infty$, then

$$K(\downarrow) \cap (L_v^\infty)_+ \neq K(\downarrow) \cap (L_w^q)_+,$$

i.e., these two cones do not coincide for any weights v, w .

If $q = \infty$, then the identity

$$K(\downarrow) \cap (L_v^\infty)_+ = K(\downarrow) \cap (L_w^\infty)_+$$

is fulfilled if and only if

$$(40) \quad \sup_{t>0} \frac{\tilde{w}(t)}{\tilde{v}(t)} < \infty, \quad \sup_{t>0} \frac{\tilde{v}(t)}{\tilde{w}(t)} < \infty.$$

Proof. Theorem 5 shows that the cones $K(\downarrow) \cap L_v^\infty$ and $K(\downarrow) \cap L_w^\infty$ coincide.

First, let $q < \infty$. Then for all $\tau \in R_+$ the function $\chi_{[0,\tau]} \frac{1}{\tilde{v}(t)}$ belongs to $K(\downarrow)$, and for its norm we have

$$(41) \quad \left\| \chi_{[0,\tau]} \frac{1}{\tilde{v}} \middle| L_v^\infty \right\| = 1, \quad \left\| \chi_{[0,\tau]} \frac{1}{\tilde{v}} \middle| L_w^q \right\| = \left(\int_0^\tau \left(\frac{w(t)}{\tilde{v}(t)} \right)^q dt \right)^{1/q}.$$

Letting τ tend to zero, we deduce from (41) that the norms $\| \cdot \|_{L_v^\infty}$ and $\| \cdot \|_{L_w^q}$ cannot be equivalent on the cone $K(\downarrow)$.

Now, let $q = \infty$. Applying Theorem 5 once again, we see that the cones $K(\downarrow) \cap L_w^\infty$ and $K(\downarrow) \cap L_v^\infty$ coincide.

The biggest function in $K(\downarrow)$ whose norm in L_v^∞ equals 1 is the function $\frac{1}{\tilde{v}}$, and the biggest function in $K(\downarrow)$ whose norm in L_w^∞ equals 1 is the function $\frac{1}{\tilde{w}}$. Conditions (40) precisely ensure the inclusions $\frac{1}{\tilde{v}} \in L_w^\infty$ and $\frac{1}{\tilde{w}} \in L_v^\infty$. \square

An analog of the Sawyer theorem in the case where $p = \infty$ looks like this.

Corollary 6. *Let $p = \infty$, and suppose we are given a measurable function $g: R_+ \rightarrow R_+$. Then*

$$\sup \left\{ \int_0^\infty x(t)g(t) dt : x \in K(\downarrow) \cap L_v^\infty, \|x\|_{L_v^\infty} \leq 1 \right\} = \int_0^\infty \frac{1}{\tilde{v}(s)} g(s) ds.$$

Proof. Theorem 5 shows that the cones $K(\downarrow) \cap L_v^\infty$ and $K(\downarrow) \cap L_w^\infty$ coincide. To finish the proof, it suffices to observe that the biggest function in $K(\downarrow)$ whose norm in L_v^∞ equals 1 is the function $\frac{1}{\tilde{v}}$. \square

Corollary 7. *Let $p = \infty$, $1 \leq q \leq \infty$. Then the Hardy operator is bounded as an operator*

$$H: K(\downarrow) \cap L_v^\infty \rightarrow L_w^q$$

if and only if

$$\left\| wH\left(\frac{1}{\tilde{v}}\right) \middle| L_w^q \right\| < \infty.$$

Theorem 6. *Fix $p = \infty$, and let X be a Banach ideal space in $S(\mu)$. For $T \in \text{Sub}(\downarrow)$ to act boundedly as an operator $T: K(\downarrow) \cap L_v^\infty \rightarrow X$, it is necessary and sufficient that*

$$\left\| T\left(\frac{1}{\tilde{v}}\right) \middle| X \right\| < \infty.$$

The two statements above are proved by much the same arguments as Corollary 6.

Now, we pass to an analog of Theorem 1 for the cone $K(\downarrow) \cap L_v^\infty$. For this, some prerequisites are needed.

To begin with, we observe that for a weight function $v: [0, \infty) \rightarrow R_+$ the condition

$$(42) \quad \limsup_{t \rightarrow \infty} v(t) = \infty$$

is equivalent to the relation

$$(43) \quad \lim_{t \rightarrow \infty} \tilde{v}(t) = \infty.$$

Note that condition (42) or the equivalent condition (43) are quite natural, because any function representable in the form $y(t) = Qx(t)$ with $x \in \bigcup_{n \in \mathbb{N}} L^1(\frac{1}{n}, \infty)$ satisfies the relation

$$(44) \quad \lim_{t \rightarrow \infty} y(t) = 0.$$

At the same time, precisely condition (43) is necessary and sufficient for an arbitrary function $x \in K(\downarrow) \cap L_v^\infty$ to satisfy $\lim_{t \rightarrow \infty} x(t) = 0$.

Theorem 7. Fix a weight function v satisfying (42) and use (34) to define a function \tilde{v} . Let there exist an absolutely continuous function $\tilde{v}_{ac} \in K(\downarrow) \cap L_v^\infty$ such that for some constant $c > 0$ we have

$$(45) \quad \frac{1}{c} \tilde{v}_{ac}(t) \leq \tilde{v}(t) \leq c \tilde{v}_{ac}(t)$$

for all $t \in \mathbb{R}_+$.

Put

$$(46) \quad \frac{1}{\tilde{v}(t)} = -\frac{d}{dt} \frac{1}{\tilde{v}_{ac}(t)}.$$

Then

(a) $Q((L_v^\infty)_+) \subset K(\downarrow) \cap L_v^\infty$, i.e., for every $x \in (L_v^\infty)_+$ we have

$$(47) \quad \|Qx \mid L_v^\infty\| \leq C \|x \mid L_v^\infty\|;$$

(b) for every $x \in K(\downarrow) \cap (L_v^\infty)_+$ there exists $x_\varepsilon \in (L_v^\infty)_+$ such that

$$\|x_\varepsilon \mid L_v^\infty\| = \|x \mid L_v^\infty\| \quad \text{and} \quad Q(x_\varepsilon)(t) \geq \frac{1}{8} x(t) \quad \text{for a.e. } t > 0.$$

The proof of Theorem 7 will be given in the last section of the paper. Also in that section, in Lemma 7, we shall indicate conditions on \tilde{v} necessary and sufficient for the existence of $\tilde{v}_{ac} \in K(\downarrow) \cap L_v^\infty$ satisfying (45). Essentially, $\tilde{v}_{ac} \in K(\downarrow) \cap L_v^\infty$ exists if and only if \tilde{v} satisfies a Δ_2 -condition at the discontinuity points.

Theorems 1 and 7 justify the following definition.

Definition 3. Suppose we are given two ideal spaces X_0, X_1 , two cones $K_0 \cap X_0, K_1 \cap X_1$, and a sublinear operator $T: K_0 \cap X_0 \rightarrow K_1 \cap X_1$. The pair $\{K_0 \cap X_0, T\}$ is said to generate the cone $K_1 \cap X_1$ if the following conditions are fulfilled:

a) there is a constant $c_0 > 0$ with the property that $\|Tx \mid X_1\| \leq c_0 \|x \mid X_0\|$ for every $x \in K_0 \cap X_0$;

b) there is a constant $c_1 > 0$ with the property that for every $y \in K_1 \cap X_1$ there exists $x_y \in K_0 \cap X_0$ such that the norm inequality $\|y \mid X_1\| \leq c_1 \|x_y \mid X_0\|$ and the a.e. inequality $y(t) \leq c_1 T x_y(t)$ both hold true.

Theorem 8. Suppose that $1 \leq p < \infty$ and a weight function v satisfying (10) is given. Define \bar{v} by (12).

Then the pair $((L_w^p)_+, Q)$ generates the cone $K(\downarrow) \cap L_v^p$ if and only if the following conditions are fulfilled:

a) the weight v satisfies (11);

b) for every $t > 0$ we have

$$(48) \quad C_7^{-1} \leq \|v \chi_{[0,t]} \mid L^p\| \cdot \left\| \frac{1}{w} \chi_{[t,\infty)} \mid L^{p'} \right\| \leq C_7$$

with a constant $C_7 > 0$ independent of t ;

c) for every $t > 0$ we have

$$(49) \quad C_8^{-1} \left\| \frac{1}{v} \chi_{[t, \infty)} \mid L^{p'} \right\| \leq \left\| \frac{1}{w} \chi_{[t, \infty)} \mid L^{p'} \right\| \leq C_8 \left\| \frac{1}{v} \chi_{[t, \infty)} \mid L^{p'} \right\|$$

with a constant $C_8 > 0$ independent of t .

Proof. First, we observe that conditions (48) and (49) are equivalent by (12).

We check the “only if” part. Suppose that the pair $((L_w^p)_+, Q)$ generates the cone $K(\downarrow) \cap L_v^p$. This implies immediately that the function $x(t) \in K(\downarrow) \cap L_v^p$ satisfies the condition

$$(50) \quad \lim_{t \rightarrow \infty} x(t) = 0.$$

For the nondegenerate cone $K(\downarrow) \cap L_v^p$, condition (12) is equivalent to the statement that the characteristic function of the entire half-line R_+ does not belong to $K(\downarrow) \cap L_v^p$. Therefore, (50) implies the necessity of (12).

We show the necessity of (48), (49). Since the pair $((L_w^p)_+, Q)$ generates the cone $K(\downarrow) \cap L_v^p$, we see that for every $t > 0$ the inequalities

$$(51) \quad \frac{1}{c_0} \|\chi_{(0, t)} \mid L_v^p\| \leq \inf \left\{ \|y \mid L_w^p\| : 1 \leq \int_t^\infty y(s) ds \right\} \leq c_0 \|\chi_{(0, t)} \mid L_v^p\|.$$

must be true with a constant $c_0 > 0$ independent of t .

By duality and formula (1), we obtain

$$1 \leq \int_t^\infty y(s) ds \leq \|y \chi_{(t, \infty)} \mid L_w^p\| \left\| \chi_{(t, \infty)} \frac{1}{w} \mid L^{p'} \right\|,$$

or

$$(52) \quad \|y \chi_{(t, \infty)} \mid L_w^p\| \geq \frac{1}{\left\| \chi_{(t, \infty)} \frac{1}{w} \mid L^{p'} \right\|}.$$

The definition of the dual space shows that we can choose a sequence y_n of functions in the unit sphere of L_w^p such that

$$\begin{aligned} & \|y_n \chi_{(t, \infty)} \mid L_w^p\| \left\| \chi_{(t, \infty)} \frac{1}{w} \mid L^{p'} \right\| \\ & \geq \int_t^\infty y_n(s) ds \geq \|y_n \chi_{(t, \infty)} \mid L_w^p\| \left\| \chi_{(t, \infty)} \frac{1}{w} \mid L^{p'} \right\| - \frac{1}{n} \\ & = \left\| \chi_{(t, \infty)} \frac{1}{w} \mid L^{p'} \right\| - \frac{1}{n}. \end{aligned}$$

Now, we define a sequence of functions z_n as follows:

$$z_n(t) = \frac{y_n(t)}{\int_t^\infty y_n(s) ds}.$$

Then

$$(53) \quad \int_t^\infty z_n(s) ds = 1 \text{ and } \|z_n \chi_{(t, \infty)} \mid L_w^p\| \leq \frac{1}{\left\| \chi_{(t, \infty)} \frac{1}{w} \mid L^{p'} \right\| - \frac{1}{n}}.$$

From (52)–(53) we deduce that

$$(54) \quad \inf \left\{ \|y \mid L_w^p\| : 1 \leq \int_t^\infty y(s) ds \right\} = \frac{1}{\left\| \chi_{(t, \infty)} \frac{1}{w} \mid L^{p'} \right\|}.$$

Conditions (51) and (54) proof the necessity of (48)–(49).

To prove that (a), (b), and (c) suffice, we may repeat the proof of Theorem 1 word-for-word. \square

Theorem 9. Suppose $p = \infty$ and we are given a weight function w . Define a function \tilde{w} by (34).

Then the pair $((L_u^\infty)_+, Q)$ generates the cone $K(\downarrow) \cap L_w^\infty$ if and only if the following relations are fulfilled:

(a) for w , we have

$$(55) \quad \limsup_{t \rightarrow \infty} w(t) = \infty;$$

(b) there exists an absolutely continuous function $\tilde{w}_{ac} \in K(\downarrow) \cap L_w^\infty$ such that, with some constant C , we have

$$(56) \quad \frac{1}{c} \tilde{w}_{ac}(t) \leq \tilde{w}(t) \leq c \tilde{w}_{ac}(t)$$

for all $t \in R_+$;

(c) for every $t > 0$ we have

$$(57) \quad C_7^{-1} \frac{1}{\tilde{w}(t)} \leq \int_t^\infty \frac{1}{u(s)} ds \leq C_7 \frac{1}{\tilde{w}(t)}$$

with a constant $C_7 > 0$ independent of t .

Proof. Theorem 5 shows that the cones $K(\downarrow) \cap L_w^\infty$ and $K(\downarrow) \cap L_{\tilde{w}}^\infty$ coincide.

We check the “only if” part. Suppose that the pair $((L_u^\infty)_+, Q)$ generates the cone $K(\downarrow) \cap L_w^\infty$. This readily implies that every function $x(t)$ in $K(\downarrow) \cap L_w^\infty$ satisfies the condition $\lim_{t \rightarrow \infty} x(t) = 0$. The last is equivalent to

$$(58) \quad \lim_{t \rightarrow \infty} \tilde{w}(t) = \infty,$$

and, as it was indicated in the proof of the equivalence of (42) and (43), condition (58) is equivalent to (55).

Next, since the pair $((L_u^\infty)_+, Q)$ generates the cone $K(\downarrow) \cap L_w^\infty$, we see that the function $\frac{1}{w} \in K(\downarrow) \cap L_w^\infty$ satisfies the inequality

$$(59) \quad \frac{1}{\tilde{w}(t)} \leq C_{10} \int_t^\infty \frac{1}{u(s)} ds$$

with a constant C_{10} independent of t . However, by Lemma 2, the boundedness condition for the operator $Q: L_u^\infty \rightarrow L_{\tilde{w}}^\infty$ looks like this:

$$(60) \quad \sup_{t \in R_+} \tilde{w}(t) \int_t^\infty \frac{1}{u(s)} ds \leq C_{11} < \infty.$$

Relations (59)–(60) prove the necessity of (b) and (c).

To prove that (a), (b), and (c) suffice, we may repeat the proof of Theorem 7 word-for-word. \square

§4. EXTRAPOLATION THEOREMS FOR CONES

In the theory of integral operators with positive kernel, a special role is played by the so-called Schur theorem or Schur test (see [25, p. 37] and [42, p. 42]), which says that an integral operator $Kx(t) = \int k(t, s)x(s) ds$ with positive kernel $k(t, s) \geq 0$ is bounded in L^p for $1 < p < \infty$ if and only if there exists a positive function u such that

$$Ku^{p'}(t) \leq Cu^{p'}(t) \quad \text{and} \quad K'u^p(t) \leq Cu^p(t),$$

here K' is the formally adjoint operator and $1/p' + 1/p = 1$. This statement can be regarded as a factorization or extrapolation theorem: there exists a positive function u (a weight function u) such that K is bounded in the following couples of spaces:

$$K: L_{u^{-p'}}^\infty \rightarrow L_{u^{-p'}}^\infty, \quad K: L_{u^p}^1 \rightarrow L_{u^p}^1.$$

We start with a reformulation of the Schur extrapolation theorem in modern terms, see [9, Corollary 7].

Proposition 1 (Schur test). *Suppose that $T, T' \in \text{Sub}(+)$, $1 < p < \infty$, and we are given two weight functions v, w . The following conditions are equivalent:*

- (a) *the operator $T: L_v^p \rightarrow L_w^p$ is bounded;*
- (b) *there exist four weight functions v_0, v_1, w_0, w_1 such that*

$$(61) \quad v(t) = v_0(t)^{1/p} v_1(t)^{1-1/p}, \quad w(t) = w_0(t)^{1/p} w_1(t)^{1-1/p}$$

for all $t \in R_+$, and the operator T acts boundedly in the following couples:

$$(62) \quad T: L_{v_0}^1 \rightarrow L_{w_0}^1, \quad T: L_{v_1}^\infty \rightarrow L_{w_1}^\infty.$$

The implication (b) \implies (a) follows from interpolation theorems for positive operators for the Calderón–Lozanovskii construction $X_0^\theta X_1^{1-\theta}$ (see [2, 26, 30, 37], [31, Theorem 15.13]) and the relations

$$\begin{aligned} (L_{v_0}^1)^{1/p} (L_{v_1}^\infty)^{1-1/p} &= L_{v_0^{1/p} v_1^{1-1/p}}^p = L_v^p, \\ (L_{w_0}^1)^{1/p} (L_{w_1}^\infty)^{1-1/p} &= L_{w_0^{1/p} w_1^{1-1/p}}^p = L_w^p. \end{aligned}$$

The reverse implication (a) \implies (b), which is the essence of theorems like the Schur test, was proved in [9, Corollary 7] (see also [5, p. 728], [8, Theorem 1], [7, p. 18]).

In this section we pass to extrapolation theorems for operators on cones. We begin with a general version of extrapolation theorems for operators on the cone $K(\downarrow)$.

Theorem 10. *Suppose that $T, T' \in \text{Sub}$, $1 < p < \infty$, and we are given a weight function v satisfying (10), (11). Define a new function \bar{v} by (12).*

Put $\theta = 1/p$. Then the following conditions are equivalent:

- a) *T is bounded as an operator in the following couple:*

$$T: K(\downarrow) \cap L_v^p \rightarrow L_u^p;$$

- b) *there exist functions v_0, v_1, u_0, u_1 satisfying*

$$(63) \quad v_0^\theta(t) \cdot v_1^{1-\theta}(t) \equiv \bar{v}(t), \quad u_0^\theta(t) \cdot u_1^{1-\theta}(t) \equiv u(t)$$

and such that TQ acts boundedly in the couples

$$(64) \quad TQ: L_{v_0}^1 \rightarrow L_{u_0}^1, \quad TQ: L_{v_1}^\infty \rightarrow L_{u_1}^\infty.$$

Proof. Suppose a) is fulfilled. Then Theorem 4 shows that this statement is equivalent to the boundedness of TQ in the couple $TQ: L_v^p \rightarrow L_u^p$. Since TQ and $(TQ)'$ belong to $\text{Sub}(+)$, we may apply the Schur test to the operator $TQ: L_v^p \rightarrow L_u^p$. This yields the conditions of item b).

Suppose b) is fulfilled. Then, by the interpolation theorem for the operator TQ , we see that TQ is bounded as indicated below:

$$TQ: (L_{v_0}^1)^\theta (L_{v_1}^\infty)^{1-\theta} \rightarrow (L_{u_0}^1)^\theta (L_{u_1}^\infty)^{1-\theta}.$$

The well-known identity

$$(L_{w_0}^1)^\theta (L_{w_1}^\infty)^{1-\theta} = L_w^p,$$

where $w_\theta(t) \equiv w_0^\theta(t) w_1^{1-\theta}(t)$, combined with (67) implies that TQ is bounded also in the following way:

$$TQ: L_{v_\theta}^p \rightarrow L_{u_\theta}^p,$$

where $v_\theta(t) \equiv v_0^\theta(t) v_1^{1-\theta}(t)$ and $u_\theta(t) \equiv u_0^\theta(t) u_1^{1-\theta}(t)$. Since $v_\theta(t) \equiv \bar{v}(t)$ and $u_\theta(t) \equiv u(t)$, relation (65) is equivalent to the boundedness of TQ in the following couple:

$$TQ: L_{\bar{v}}^p \rightarrow L_u^p.$$

But $T \in \text{Sub}(+)$, and, by Theorem 8, the pair $((L_v^p)_+, Q)$ generates the cone $K(\downarrow) \cap L_v^p$; therefore, the last relation implies a). \square

Theorem 10 has an important drawback. It would be desirable to replace conditions (63) and (64) by the following more natural conditions:

there exist functions v_0, v_1, u_0, u_1 satisfying

$$(65) \quad v_0^\theta(t) \cdot v_1^{1-\theta}(t) \equiv v(t), \quad u_0^\theta(t) \cdot u_1^{1-\theta}(t) \equiv u(t)$$

and such that T acts boundedly in the following couples:

$$(66) \quad T: K(\downarrow) \cap L_{v_0}^1 \rightarrow L_{u_0}^1, \quad T: K(\downarrow) \cap L_{v_1}^\infty \rightarrow L_{u_1}^\infty.$$

We are going to obtain an analog of Theorem 10 with conditions (63), (64) replaced by (65), (66) in one important particular case. Some preliminaries are required for this.

Let X_0, X_1 be two ideal spaces with $X_0, X_1 \subset S(\mu)$. Fix $0 < \theta < 1$. The new ideal space $X_0^\theta X_1^{1-\theta}$ (the Calderón–Lozanovskii construction) consists of all $x \in S(\mu)$ for which the following norm is finite:

$$(67) \quad \|x \mid X_0^\theta X_1^{1-\theta}\| = \inf \{ \lambda > 0 : |x(t)| \leq \lambda \cdot |x_0(t)|^\theta |x_1(t)|^{1-\theta} \forall t \in [0, \infty); \\ \|x_0 \mid X_0\| \leq 1, \quad \|x_1 \mid X_1\| \leq 1 \}.$$

The space $X_0^\theta X_1^{1-\theta}$ was introduced by Calderón in [13] for the study of the complex interpolation method.

Definition 4. A cone K is said to be *canonical* if for every pair x, y of functions in K and every number $\theta \in (0, 1)$ the function $x^\theta \cdot y^{1-\theta}$ again belongs to K .

We observe that the cones of monotonic functions are canonical.

If K is a canonical cone in $S(\mu)$, then by analogy with the space $X_0^\theta X_1^{1-\theta}$ we can introduce the new cone $(K \cap X_0)^\theta (K \cap X_1)^{1-\theta}$, admitting in (67) only decompositions that involve elements of the cone.

Remark 1. It is easily seen that for a canonical cone and $\theta \in (0, 1)$ we always have a continuous embedding

$$(K \cap X_0)^\theta (K \cap X_1)^{1-\theta} \subseteq K \cap X_0^\theta X_1^{1-\theta}.$$

On the other hand, as it usually happens in interpolation theory, for an arbitrary canonical cone K the relation

$$(K \cap X_0)^\theta (K \cap X_1)^{1-\theta} = K \cap X_0^\theta X_1^{1-\theta}$$

may fail. Even for the best studied cone $K(\downarrow)$, no sharp conditions are known that ensure this relation in the scale of Lebesgue spaces.

The next theorem is of interpolation nature. It is well known for the cone consisting of nonnegative functions (see, e.g., [2, 30, 31]).

Theorem 11. Suppose that T is a positive operator and K_0, K_1 are two canonical cones in $S(\mu)_+$. Consider four Banach ideal spaces X_0, X_1, Y_0, Y_1 in $S(\mu)$ and suppose that T acts boundedly as indicated: $T: K_0 \cap X_i \rightarrow K_1 \cap Y_i$, ($i = 0, 1$). Fix $\theta \in (0, 1)$. Then for every $x_0 \in K_0 \cap X_0$, $x_1 \in K_1 \cap X_1$ we have the pointwise inequality

$$(68) \quad T(x_0^\theta \cdot x_1^{1-\theta})(t) \leq (Tx_0(t))^\theta \cdot (Tx_1(t))^{1-\theta},$$

and T acts boundedly when regarded as an operator $T: (K_0 \cap X_0)^\theta (K_0 \cap X_1)^{1-\theta} \rightarrow (K_1 \cap Y_0)^\theta (K_1 \cap Y_1)^{1-\theta}$.

Proof. Take $x_0 \in K_0 \cap X_0$, $x_1 \in K_0 \cap X_1$ and construct the element $x_0^\theta \cdot x_1^{1-\theta} \in (K \cap X_0)^\theta (K \cap X_1)^{1-\theta}$. The numerical identity

$$(69) \quad a^\theta \cdot b^{1-\theta} = \inf_{\varepsilon > 0} \{ \varepsilon \theta a + \varepsilon^{-\frac{\theta}{1-\theta}} (1-\theta) b \},$$

valid for all $a > 0$, $b > 0$, implies the inequality

$$(70) \quad \begin{aligned} T(x_0^\theta \cdot x_1^{1-\theta})(t) &\leq T(\varepsilon \theta x_0(t) + \varepsilon^{-\frac{\theta}{1-\theta}} (1-\theta) x_1(t)) \\ &\leq \varepsilon \theta T x_0(t) + \varepsilon^{-\frac{\theta}{1-\theta}} (1-\theta) T x_1(t). \end{aligned}$$

Minimizing the right-hand side in (70) over $\varepsilon > 0$ for each fixed t and taking (69) into account, we arrive at (68).

The boundedness of T in the required sense,

$$T: (K_0 \cap X_0)^\theta (K_0 \cap X_1)^{1-\theta} \rightarrow (K_1 \cap Y_0)^\theta (K_1 \cap Y_1)^{1-\theta},$$

is immediate from (68). \square

Lemma 3. Fix $\theta \in (0, 1)$. In the spaces $L_{w_0}^1$, $L_{w_1}^\infty$, and $L_{u_0}^1$, consider the cones $K(\downarrow) \cap L_{w_0}^1$, $K(\downarrow) \cap L_{w_1}^\infty$, and $K(\downarrow) \cap L_{u_0}^1$ for which we have a continuous embedding

$$(K(\downarrow) \cap L_{w_0}^1)^\theta (K(\downarrow) \cap L_{w_1}^\infty)^{1-\theta} \subseteq (K(\downarrow) \cap L_{u_0}^1)^\theta (K(\downarrow) \cap L_{w_1}^\infty)^{1-\theta}.$$

Then we have a continuous embedding

$$K(\downarrow) \cap L_{w_0}^1 \subseteq K(\downarrow) \cap L_{u_0}^1.$$

Proof. Lemma 8 in the last section shows that for every z of unit norm in $(K(\downarrow) \cap L_{w_0}^1)^\theta (K(\downarrow) \cap L_{w_1}^\infty)^{1-\theta}$ we have

$$(71) \quad z(t) \leq x_0^\theta(t) \cdot \left(\frac{1}{\tilde{w}(t)} \right)^{1-\theta} (t) \quad (t \in R_+)$$

with $x_0 \in K(\downarrow) \cap L_{w_0}^1$ and $\|x_0\|_{L_{w_0}^1} = 1$.

Since \tilde{w} is nonzero a.e., the claim follows from (71). \square

Now everything is ready to prove an analog of Theorem 10 with conditions (63) and (64) replaced by (65) and (66).

Theorem 12. Fix $p \in (1, \infty)$ and a weight function v satisfying (10), (11). Define a new function \bar{v} by (12) and put $\theta = 1/p$.

Suppose we are given operators $T, T' \in \text{Sub}(+)$, where $T \in \text{Sub}(+)$ acts boundedly in the couple

$$T: K(\downarrow) \cap L_v^p \rightarrow L_v^p.$$

Then there exist functions w_0, w_1 such that

$$w_0^\theta(t) \cdot w_1^{1-\theta}(t) \equiv v(t),$$

and T acts boundedly in the following couples:

$$T: K(\downarrow) \cap L_{w_0}^1 \rightarrow L_{w_0}^1, \quad T: K(\downarrow) \cap L_{w_1}^\infty \rightarrow L_{w_1}^\infty.$$

Proof. We introduce a new operator T_1 by the formula

$$T_1 x(t) = T x(t) + x(t)$$

and apply Theorem 10 to it, obtaining functions v_0, v_1, w_0, w_1 for which we have

$$(72) \quad v_0^\theta(t) \cdot v_1^{1-\theta}(t) \equiv \bar{v}(t), \quad w_0^\theta(t) \cdot w_1^{1-\theta}(t) \equiv v(t)$$

(here \bar{v} is defined by (12)).

The operator $T_1 Q$ acts boundedly in the following couples:

$$TQ + Q: L_{v_0}^1 \rightarrow L_{w_0}^1, \quad TQ + Q: L_{v_1}^\infty \rightarrow L_{w_1}^\infty.$$

We introduce a function u_0 by the formula

$$(73) \quad \|\chi_{(0,t)} u_0\|_{L^1} \cdot \left\| \chi_{(t,\infty)} \frac{1}{v_0} \right\|_{L^\infty} \equiv 1$$

and consider the space $L_{u_0}^1$. By (73) and Theorem 8, the pair $((L_{v_0}^1)_+, Q)$ generates the cone $K(\downarrow) \cap L_{u_0}^1$.

Next, we introduce a function u_1 by the formula

$$(74) \quad \frac{1}{u_1(t)} \equiv \int_t^\infty \frac{1}{v_1(s)} ds$$

and consider the space $L_{u_1}^\infty$. By (74) and Theorem 9, the pair $((L_{v_1}^\infty)_+, Q)$ generates the cone $K(\downarrow) \cap L_{u_1}^\infty$.

Since Q acts boundedly in the couples

$$Q: L_{v_0}^1 \rightarrow L_{w_0}^1, \quad Q: L_{v_1}^\infty \rightarrow L_{w_1}^\infty$$

and the cones $K(\downarrow) \cap L_{u_0}^1$, $K(\downarrow) \cap L_{u_1}^\infty$ are generated by the pairs $((L_{v_0}^1)_+, Q)$ and $((L_{v_1}^\infty)_+, Q)$, respectively, we arrive at the continuous embeddings

$$(75) \quad K(\downarrow) \cap L_{u_0}^1 \subseteq K(\downarrow) \cap L_{w_0}^1, \quad K(\downarrow) \cap L_{u_1}^\infty \subseteq K(\downarrow) \cap L_{w_1}^\infty.$$

On the other hand, since Q is positive, inequality (68) in Theorem 11 shows that for every $x_0 \in L_{v_0}^1$, $x_1 \in L_{v_1}^\infty$ we have

$$(76) \quad Q(x_0^\theta \cdot x_1^{1-\theta})(t) \leq (Qx_0(t))^\theta \cdot (Qx_1(t))^{1-\theta}$$

almost everywhere. By Theorem 8, the pair $((L_v^p)_+, Q)$ generates the cone $K(\downarrow) \cap L_v^p$. Therefore, (76) yields the continuous embedding

$$(77) \quad K(\downarrow) \cap L_v^p \subseteq (K(\downarrow) \cap L_{u_0}^1)^\theta (K(\downarrow) \cap L_{u_1}^\infty)^{1-\theta}.$$

At the same time, formula (72), Remark 1, and the definitions yield the continuous embedding

$$(78) \quad (K(\downarrow) \cap L_{w_0}^1)^\theta (K(\downarrow) \cap L_{w_1}^\infty)^{1-\theta} \subseteq K(\downarrow) \cap L_v^p.$$

Consequently, by (77) and (78) we obtain

$$(K(\downarrow) \cap L_{w_0}^1)^\theta (K(\downarrow) \cap L_{w_1}^\infty)^{1-\theta} \subseteq K(\downarrow) \cap L_v^p \subseteq (K(\downarrow) \cap L_{u_0}^1)^\theta (K(\downarrow) \cap L_{u_1}^\infty)^{1-\theta}.$$

This implies the continuous embedding

$$(K(\downarrow) \cap L_{w_0}^1)^\theta (K(\downarrow) \cap L_{w_1}^\infty)^{1-\theta} \subseteq (K(\downarrow) \cap L_{u_0}^1)^\theta (K(\downarrow) \cap L_{u_1}^\infty)^{1-\theta}.$$

Together with the second embedding in (75), this yields the continuity of the embeddings

$$\begin{aligned} (K(\downarrow) \cap L_{w_0}^1)^\theta (K(\downarrow) \cap L_{u_1}^\infty)^{1-\theta} &\subseteq (K(\downarrow) \cap L_{w_0}^1)^\theta (K(\downarrow) \cap L_{w_1}^\infty)^{1-\theta} \\ &\subseteq (K(\downarrow) \cap L_{u_0}^1)^\theta (K(\downarrow) \cap L_{u_1}^\infty)^{1-\theta}. \end{aligned}$$

Thus,

$$(79) \quad (K(\downarrow) \cap L_{w_0}^1)^\theta (K(\downarrow) \cap L_{u_1}^\infty)^{1-\theta} \subseteq (K(\downarrow) \cap L_{u_0}^1)^\theta (K(\downarrow) \cap L_{u_1}^\infty)^{1-\theta}.$$

By (79), using Lemma 3, we obtain the embedding

$$K(\downarrow) \cap L_{w_0}^1 \subseteq K(\downarrow) \cap L_{u_0}^1.$$

Comparing this with the first embedding in (75), we see that, up to equivalent norms, we have

$$(80) \quad K(\downarrow) \cap L_{w_0}^1 = K(\downarrow) \cap L_{u_0}^1.$$

Thus, (73) and (80) imply the existence of a constant $c > 0$ such that for every $t > 0$ we have

$$c < \|\chi_{(0,t)} w_0\|_{L^1} \cdot \left\| \chi_{(t,\infty)} \frac{1}{v_0} \right\|_{L^\infty} < \frac{1}{c}.$$

Combining (12) and (72) with the last relation, we see that

$$\begin{aligned} 1 &= \left(\int_0^t (w_0^\theta(s) \cdot w_1^{1-\theta}(s))^{1/\theta} ds \right)^\theta \cdot \left(\int_t^\infty \left(\frac{1}{v_0^\theta(s) \cdot v_1^{1-\theta}(s)} \right)^{1/(1-\theta)} ds \right)^{1-\theta} \\ &= \left(\int_0^t w_0(s) \cdot w_1^{(1-\theta)/\theta}(s) ds \right)^\theta \cdot \left(\int_t^\infty \left(\frac{1}{v_0(s)} \right)^{\theta/(1-\theta)} \left(\frac{1}{v_1(s)} \right) ds \right)^{1-\theta} \\ &\leq \sup_{s \leq t} w_1^{1-\theta}(s) \cdot \left(\int_0^t w_0(s) ds \right)^\theta \cdot \sup_{s \geq t} \frac{1}{v_0^\theta(s)} \cdot \left(\int_t^\infty \frac{1}{v_1(s)} ds \right)^{1-\theta} \\ &= \left(\sup_{s \leq t} w_1(s) \cdot \int_t^\infty \frac{1}{v_1(s)} ds \right)^{1-\theta} \cdot \left(\sup_{s \geq t} \frac{1}{v_0(s)} \cdot \int_0^t w_0(s) ds \right)^\theta \\ &\leq \left(\frac{1}{c} \right)^\theta \cdot \left(\sup_{s \leq t} w_1(s) \cdot \int_t^\infty \frac{1}{v_1(s)} ds \right)^{1-\theta} \end{aligned}$$

for all $t \in R_+$. This shows that the inequality

$$c^{\frac{\theta}{1-\theta}} \cdot \inf_{s \leq t} \frac{1}{w_1(s)} \leq \int_t^\infty \frac{1}{v_1(s)} ds$$

is true for all $t \in R_+$. Taking (74) and (34) into account, we can rewrite the last inequality in the following equivalent form: for all $t \in R_+$ we have

$$\frac{1}{\tilde{w}_1(t)} \leq c^{\frac{-\theta}{1-\theta}} \frac{1}{u_1(t)},$$

or, in the language of embeddings,

$$(81) \quad K(\downarrow) \cap L_{\tilde{w}_1}^\infty \subseteq K(\downarrow) \cap L_{u_1}^\infty.$$

Theorem 5 implies the identity $K(\downarrow) \cap L_{w_1}^\infty = K(\downarrow) \cap L_{\tilde{w}_1}^\infty$. Thus, the embedding (81) is equivalent to the embedding

$$K(\downarrow) \cap L_{w_1}^\infty \subseteq K(\downarrow) \cap L_{\tilde{u}_1}^\infty.$$

Together with the second embedding in (75), this relation shows that, up to equivalent norms, the following identity holds true:

$$K(\downarrow) \cap L_{u_1}^\infty = K(\downarrow) \cap L_{w_1}^\infty.$$

Combined with (80), this proves the theorem. \square

Since the Hardy operator fits in the scope of Theorem 12, we have the following statement.

Theorem 13. Fix $p \in (1, \infty)$ and consider a weight function v satisfying (10) and (11). Put $\theta = 1/p$. The Hardy operator H is bounded as an operator

$$H: K(\downarrow) \cap L_v^p \rightarrow L_v^p$$

if and only if there exist functions w_0, w_1 such that a) $w_0^\theta(t) \cdot w_1^{1-\theta}(t) \equiv v(t)$ for all $t \in R_+$;

b) H acts boundedly in the couples

$$H: K(\downarrow) \cap L_{w_0}^1 \rightarrow L_{w_0}^1, \quad H: K(\downarrow) \cap L_{w_1}^\infty \rightarrow L_{w_1}^\infty.$$

§5. PROOFS OF THEOREMS 1 AND 7,
AND AUXILIARY LEMMAS

To prove Theorem 1, we need some auxiliary statements, with which we shall start.

Lemma 4. *Let X be an ideal space. Take a numerical sequence*

$$0 < \cdots < t_j < t_{j+1} < \cdots < \infty \text{ with } \lim_{j \rightarrow -\infty} t_j = 0.$$

Let the element $x = \sum_{-\infty}^{\infty} 2^{-j} \chi_{[0, t_j)}$ belong to X and satisfy the condition

$$(82) \quad \lim_{k \rightarrow -\infty} \left\| \sum_{j=-\infty}^k 2^{-j} \chi_{[0, t_j)} \mid X \right\| = 0.$$

Then there exists a sequence of integers $k_j : -\infty < \cdots < k_j < k_{j-1} < \cdots < k_0 < \infty$ such that

$$\sum_{i=0}^{\infty} \left\| \sum_{j=k_{i+1}+1}^{k_i} 2^{-j} \chi_{[0, t_j)} \mid X \right\| + \left\| \sum_{j=k_0}^{\infty} 2^{-j} \chi_{[0, t_j)} \mid X \right\| \leq 2 \|x \mid X\|.$$

Proof. The sequence k_i can be defined as follows. Using (82), take k_0 so as to have

$$\left\| \sum_{-\infty}^{k_0} 2^{-j} \chi_{[0, t_j)} \mid X \right\| \leq 2^{-1} \|x \mid X\|.$$

Suppose that the numbers $k_{i-1} < k_{i-2} < \cdots < k_1 < k_0$ are constructed. Then we choose $k_i < k_{i-1}$ so as to have

$$\left\| \sum_{-\infty}^{k_i} 2^{-j} \chi_{[0, t_j)} \mid X \right\| \leq 2^{-i-1} \|x \mid X\|.$$

The possibility of this choice follows from (82).

Since X is an ideal space, an easy calculation shows that

$$\begin{aligned} & \sum_{i=0}^{\infty} \left\| \sum_{j=k_{i+1}+1}^{k_i} 2^{-j} \chi_{[0, t_j)} \mid X \right\| + \left\| \sum_{j=k_0+1}^{\infty} 2^{-j} \chi_{[0, t_j)} \mid X \right\| \\ & \leq \sum_{i=0}^{\infty} \left\| \sum_{-\infty}^{k_i} 2^{-j} \chi_{[0, t_j)} \mid X \right\| + \left\| \sum_{j=k_0+1}^{\infty} 2^{-j} \chi_{[0, t_{j+1})} \mid X \right\| \\ & \leq \sum_{i=0}^{\infty} 2^{-i-1} \|x \mid X\| + \|x \mid X\| \leq 2 \|x \mid X\|. \quad \square \end{aligned}$$

The next lemma allows us to estimate the norms of certain specific functions.

Lemma 5. *Let $p \in [1, \infty)$. Consider the space L_v^p , where the weight v satisfies (10) for every $t > 0$, i.e., the cone $K(\downarrow) \cap L_v^p$ is nondegenerate.*

Let $\{t_j\}_{j=k}^{\infty}$ be a numerical sequence such that the relations $t_{j+1} > t_j$ are fulfilled for all $j = k, k+1, \dots$. Suppose that the element $x = \sum_{j=k}^{\infty} 2^{-j} \chi_{[0, t_j)}$ belongs to L_v^p . Then we have

$$(83) \quad \|x \mid L_v^p\|^p = 2^{(1-k)p} \|\chi_{[0, t_k)} v \mid L^p\|^p + \sum_{j=k}^{\infty} 2^{-(j+1)p} \|\chi_{[t_j, t_{j+1})} v \mid L^p\|^p,$$

$$(84) \quad \sum_{j=k}^{\infty} 2^{-jp} \|\chi_{[0,t_j]} v \mid L^p\|^p = \frac{1}{1-2^{-p}} (2^{-kp} \|\chi_{[0,t_k]} v \mid L^p\|^p + \sum_{j=k}^{\infty} 2^{-(j+1)p} \|\chi_{[t_j,t_{j+1}]} v \mid L^p\|^p).$$

Proof. First, we verify (83):

$$\begin{aligned} \|x \mid L_v^p\|^p &= \left\| \sum_{j=k}^{\infty} 2^{-j} \chi_{[0,t_j]} \mid L_v^p \right\|^p = \left\| 2^{-k+1} \chi_{[0,t_k]} + \sum_{j=k}^{\infty} 2^{-j} \chi_{[t_j,t_{j+1}]} \mid L_v^p \right\|^p \\ &= 2^p \|2^{-k} \chi_{[0,t_k]} \mid L_v^p\|^p + \sum_{j=k}^{\infty} 2^{-pj} \|\chi_{[t_j,t_{j+1}]} \mid L_v^p\|^p. \end{aligned}$$

Next, we prove (84). Condition (10) shows that for every t_j we have $\|\chi_{[0,t_j]} v \mid L^p\|^p < \infty$. Therefore, for every $m \geq k$ we have

$$(85) \quad \|\chi_{[0,t_m]} v \mid L^p\|^p = \|\chi_{[0,t_k]} v \mid L^p\|^p + \sum_{i=k}^{m-1} \|\chi_{[t_i,t_{i+1}]} v \mid L^p\|^p.$$

Suppose first that the left-hand side in (84) is finite. Then, taking (85) into account, we obtain

$$(86) \quad \begin{aligned} &\sum_{j=k}^{\infty} 2^{-jp} \|\chi_{[0,t_j]} v \mid L^p\|^p \\ &= \|\chi_{[0,t_k]} v \mid L^p\|^p \sum_{j=k}^{\infty} 2^{-jp} + \sum_{i=k}^{\infty} \left(\sum_{j=i+1}^{\infty} 2^{-jp} \right) \|\chi_{[t_i,t_{i+1}]} v \mid L^p\|^p \\ &= \frac{2^{-kp}}{1-2^{-p}} \|\chi_{[0,t_k]} v \mid L^p\|^p + \sum_{i=k}^{\infty} \frac{2^{-(i+1)p}}{1-2^{-p}} \|\chi_{[t_i,t_{i+1}]} v \mid L^p\|^p. \end{aligned}$$

Thus, (84) is true in this case.

But if the left-hand side of (84) is infinite, we deduce that the right-hand side is also infinite because all transformations in (86) have been done for nonnegative terms. \square

The next statement is a principal lemma in this paper.

Lemma 6. Fix $p \in [1, \infty)$ and a weight function v such that (10) is true for all $t > 0$, i.e., the cone $K(\downarrow) \cap L_v^p$ is nondegenerate.

We introduce a new function \bar{v} by the equation

$$(87) \quad \|\chi_{[0,t]} v \mid L^p\| \cdot \left\| \chi_{[t,\infty)} \frac{1}{\bar{v}} \mid L^{p'} \right\| \equiv 1.$$

Then if a function x from the unit ball of L_v^p has the form $x = \sum_{i=k_0}^{k_1} 2^{-i} \chi_{[0,t_i]}$ ($0 \leq t_{k_0} < \dots < t_i < t_{i+1} < \dots < \infty$, where k_0 is finite and k_1 may be infinite), then for every $\varepsilon > 0$ there exists a function $x_\varepsilon \in L_{\bar{v}}^p$ satisfying

$$(88) \quad \|x_\varepsilon \mid L_{\bar{v}}^p\| \leq \frac{1+\varepsilon}{(2^p-1)^{1/p}} \|x \mid L_v^p\|$$

and such that for all $t \in [0, \infty)$ we have

$$(89) \quad (Qx_\varepsilon)(t) \geq \frac{1}{16} x(t).$$

Proof. So, suppose we are given a function x in the unit ball of L_v^p and $x = \sum_{i=k_0}^{k_1} 2^{-i} \chi_{[0, t_i)}$ ($0 \leq t_{k_0} < \dots < t_i < t_{i+1} < \dots < \infty$, k_0 is finite and k_1 may be infinite).

It is easily seen that for every admissible $i = k_0, k_0 + 1, \dots$ we have

$$(90) \quad 2^{-i-1} \leq x(t_i) \leq 2^{-i}.$$

For every admissible $i \in Z$, we define a number b_i by

$$(91) \quad b_i = \inf \left\{ \|y\|_{L_v^p} : \int_{t_i}^{\infty} y(s) ds \geq 2^{-i-1} \right\}.$$

Since $t_i < \infty$ and the weight function is finite a.e., all numbers b_i are finite. Moreover, since $p \in [1, \infty)$ and L_v^p is an ideal space, it follows that

$$(92) \quad \inf \left\{ \|y\|_{L_v^p} : \int_{t_i}^{\infty} y(s) ds \geq 2^{-i-1} \right\} = \inf \left\{ \|y\|_{L_v^p} : \int_{t_i}^{\infty} y(s) ds = 2^{-i-1} \right\},$$

i.e., we may assume that we have equality in (91). By the definition of the dual space, formulas (87) and (92) yield immediately two important relations:

$$(93) \quad \begin{aligned} 2^{-i-1} &= \int_{t_i}^{\infty} y(s) ds \leq \|\chi_{[t_i, \infty)} y\|_{L_v^p} \cdot \|\chi_{[t_i, \infty)}\|_{(L_v^p)'} \\ &= \|\chi_{[t_i, \infty)} y\|_{L_v^p} \cdot \left\| \chi_{[t_i, \infty)} \frac{1}{v} \right\|_{L^{p'}}, \end{aligned}$$

$$(94) \quad b_i = \frac{2^{-i-1}}{\|\chi_{[t_i, \infty)} \frac{1}{v}\|_{L^{p'}}} = 2^{-i-1} \cdot \|\chi_{[0, t_i)} v\|_{L^p}.$$

Fixing $\varepsilon > 0$, for every $i = k_0, k_0 + 1, \dots$ we choose a nearly extremal function y_i ensuring the relations

$$(95) \quad \text{supp } y_i \subseteq [t_i, \infty), \quad \int_{t_i}^{\infty} y_i(s) ds = 2^{-i-1}, \quad b_i \leq \|y_i\|_{L_v^p} \leq b_i(1 + \varepsilon).$$

The possibility of such a choice is clear.

The subsequent construction of the required function is entirely algorithmic. So, we present it in the form usual for description of algorithms. Thus, let a collection of functions $\{y_k(t)\}_{k_0}^{k_1}$ be given.

Fix $k_0 \in Z$. Put $k = k_0$, $\zeta_{k_0}(t) = y_{k_0}(t)$.

Step A. If

$$(96) \quad \int_{t_k}^{t_{k+1}} \zeta_k(s) ds \geq \frac{1}{2} \int_{t_k}^{\infty} \zeta_k(s) ds,$$

put $z_k(t) = \zeta_k(t) \chi_{[t_k, t_{k+1})}$, $k = k + 1$, $\zeta_k(t) = y_k(t)$. Return to Step A.

Step B. If (96) fails, i.e., we have

$$(97) \quad \int_{t_k}^{t_{k+1}} \zeta_k(s) ds < \frac{1}{2} \int_{t_k}^{\infty} \zeta_k(s) ds,$$

then again define z_k by $z_k(t) = \zeta_k(t) \chi_{[t_k, t_{k+1})}$, remove the function y_{k+1} from the collection $\{y_k(t)\}_{k_0}^{k_1}$, put $\zeta_{k+1}(t) \equiv \zeta_k(t)$, $k = k + 1$, and return to Step A.

Note that if $k_1 < \infty$, then the last step of the algorithm is done for $k = k_1 - 1$. In this case z_{k_1} should be modified. Specifically, if the last step of the algorithm is of type **B**, we define z_{k_1} by $z_{k_1}(t) = y_{k_1}(t)$, but if the last step of the algorithm is of type **B**, we put $z_{k_1}(t) = \zeta_{k_1-1} \chi_{[t_{k_1}, \infty)}$.

First, we show that for all admissible k : $k \geq k_0$ we have

$$(98) \quad \int_{t_k}^{\infty} \zeta_k(s) ds \geq \int_{t_k}^{\infty} y_k(s) ds.$$

Indeed, for $k = k_0$ we have equality in (98). We do an induction step. If we perform Step **A** after Step **A**, then again we have equality in (98). If we perform Step **A** after Step **B**, then from (97) and the inductive hypothesis we deduce that

$$\begin{aligned} \int_{t_{k+1}}^{\infty} \zeta_{k+1}(s) ds &= \int_{t_{k+1}}^{\infty} \zeta_k(s) ds = \int_{t_k}^{\infty} \zeta_k(s) ds - \int_{t_k}^{t_{k+1}} \zeta_k(s) ds \\ &\geq \frac{1}{2} \int_{t_k}^{\infty} \zeta_k(s) ds \geq \frac{1}{2} \int_{t_k}^{\infty} y_k(s) ds = \int_{t_{k+1}}^{\infty} y_{k+1}(s) ds. \end{aligned}$$

Thus, (98) is proved.

The algorithm results in replacing the collection $\{y_k(t)\}_{k_0}^{k_1}$ with a new collection of functions $\{z_k(t)\}_{k_0}^{k_1}$; furthermore, the procedure implies directly that the supports of the functions in the collection $\{z_k(t)\}_{k_0}^{k_1}$ are mutually disjoint.

Now, we define the new function

$$\overline{x_{k_0}}(t) = \sum_{k_0}^{k_1} z_k(t).$$

First, we show that for every $j = k_0, k_0 + 1, \dots$ we have

$$(99) \quad Q\overline{x_{k_0}}(t_j) = \int_{t_j}^{\infty} \sum_{k_0}^{k_1} z_k(s) ds \geq \frac{1}{4} \int_{t_j}^{\infty} y_j(s) ds = 2^{-j-3} \geq 2^{-3} x(t_j).$$

Three possibilities may occur.

a) Let $z_j(t) \equiv y_j(t)\chi_{[t_j, t_{j+1})}$, $z_{j+1}(t) \equiv y_{j+1}(t)\chi_{[t_{j+1}, t_{j+2})}$, i.e., Step **A** is performed. Then (96) yields

$$Q\overline{x_{k_0}}(t_j) = \int_{t_j}^{\infty} \sum_{k_0}^{k_1} z_k(s) ds \geq \int_{t_j}^{t_{j+1}} z_j(s) ds = \int_{t_j}^{t_{j+1}} y_j(s) ds \geq \frac{1}{2} \int_{t_j}^{\infty} y_j(s) ds,$$

which proves (99) in the case in question.

b) Suppose that, starting with some $m \geq k_0$, we have

$$z_k(t)\chi_{[t_k, t_{k+1})} \equiv y_m(t)\chi_{[t_k, t_{k+1})}, \quad k = m, m+1, \dots, m+l \quad (1 < l < \infty),$$

and that $z_{m+l+1}(t)$ does not coincide with $y_m(t)$ on $[t_{m+l+1}, t_{m+l+2})$. This happens if, starting with $k = m$, the algorithm walks away to Step **B** and does not change the function $\zeta_m(t)$ $(l-1)$ times, i.e., in accordance with (96) and (97), we have the relations

$$(100) \quad \begin{aligned} \int_{t_j}^{t_{j+1}} y_m(s) ds &< \frac{1}{2} \int_{t_{j+1}}^{\infty} y_m(s) ds \quad \text{for } j = m, m+1, \dots, m+l-1; \\ \int_{t_{m+l}}^{t_{m+l+1}} y_m(s) ds &\geq \frac{1}{2} \int_{t_{m+l+1}}^{\infty} y_m(s) ds. \end{aligned}$$

In this case, for $j = m, m+1, \dots, m+l$ we put

$$(101) \quad \begin{aligned} a_j &= \int_{t_j}^{t_{j+1}} \zeta_j(s) ds = \int_{t_j}^{t_{j+1}} z_j(s) ds = \int_{t_j}^{t_{j+1}} y_m(s) ds, \\ d_j &= \int_{t_{j+1}}^{\infty} \zeta_j(s) ds = \int_{t_{j+1}}^{\infty} y_m(s) ds. \end{aligned}$$

Therefore, (100) and (101) show that for all $j = m, m+1, \dots, m+l$ the following inequalities hold true:

$$\begin{aligned} Q\overline{x_{k_0}}(t_j) &= \int_{t_j}^{\infty} \sum_{k_0}^{k_1} z_k(s) ds \geq \int_{t_j}^{t_{m+l+1}} \sum_{i=j}^{m+l} z_j(s) ds \\ &= \int_{t_j}^{t_{j+m+l}} y_m(s) ds = a_j + a_{j+1} + \dots + a_{m+l} \\ &= a_j + a_{j+1} + \dots + a_{m+l-1} + \frac{1}{2}a_{m+l} + \frac{1}{2}a_{m+l} \\ &\geq a_j + a_{j+1} + \dots + a_{m+l-1} + \frac{1}{2}a_{m+l} + \frac{1}{4}d_{m+l} \\ &\geq \frac{1}{4} \int_{t_j}^{\infty} y_m(s) ds = \frac{1}{4} \int_{t_j}^{\infty} \zeta_j(s) ds. \end{aligned}$$

To prove (99) in the case in question, it remains to apply (98).

c) It remains to consider the case where, starting with some $m \geq k_0$, for all $k \geq m$ we have $z_k(t) \equiv y_m(t)$. This happens if, starting with $k = m$, the algorithm walks away to Step **B** and does not change the function $\zeta_m(t)$ any longer. In this case, for all $j = m, m+1, \dots$ we have $\zeta_m(t) \equiv \zeta_j(t) \equiv y_m(t)$. By (98), it follows that we also have

$$Q\overline{x_{k_0}}(t_j) = \int_{t_j}^{\infty} \sum_{k_0}^{k_1} z_k(s) ds = \int_{t_j}^{\infty} y_m(s) ds = \int_{t_j}^{\infty} \zeta_j(s) ds \geq \int_{t_j}^{\infty} y_j(s) ds.$$

Again, this implies (99).

Now, let $t \in (t_j, t_{j+1})$. Then the explicit form of the function x , inequalities (90) and the nonnegativity of $\overline{x_{k_0}}(t)$ imply the relation

$$Q\overline{x_{k_0}}(t) = \int_t^{\infty} \sum_{k_0}^{k_1} z_k(s) ds \geq \int_{t_{j+1}}^{\infty} \sum_{k_0}^{k_1} z_k(s) ds \geq 2^{-j-4} \geq \frac{1}{16}x(t_j) \geq \frac{1}{16}x(t).$$

Put $x_{\varepsilon}(t) \equiv \overline{x_{k_0}}(t)$. The last inequality and (99) yield (89). It remains to verify (88). First, we prove the relation

$$(102) \quad \|x_{\varepsilon}\|_{L_v^p}^p = \sum_{j=k_0}^{k_1} \|z_j\|_{L_v^p}^p \leq \sum_{j=k_0}^{k_1} \|y_j\|_{L_v^p}^p.$$

The identity in (102) follows from the fact that the supports of the functions in the collection $\{z_j(t)\}_{k_0}^{k_1}$ are mutually disjoint. On the other hand, for every function y_j the algorithm either merely drops it or multiplies it by a characteristic function, i.e.,

$$\sum_{j=k_0}^{k_1} z_j = \sum_{j=k_0}^{k_1} \chi(D_j)y_j,$$

where $D_j = \emptyset$ if y_j was dropped, $D_j = [t_j, t_{j+1})$ if y_j was involved in the action of the algorithm only once, $D_j = [t_j, t_{j+2})$ if y_j was involved in the action of the algorithm twice, and so on. Therefore,

$$\sum_{j=k_0}^{k_1} \|z_j\|_{L_v^p}^p = \sum_{j=k_0}^{k_1} \|y_j \chi(D_j)\|_{L_v^p}^p \leq \sum_{j=k_0}^{k_1} \|y_j\|_{L_v^p}^p.$$

So, relations (102) are proved.

Next, taking the choice of the y_j into account, by (95) and (94) we obtain the inequality

$$\|x_\varepsilon | L_v^p\|^p \leq \sum_{j=k_0}^{k_1} \|y_j | L_v^p\|^p \leq (1+\varepsilon)^p \sum_{j=k_0}^{k_1} b_j^p \leq (1+\varepsilon)^p \sum_{j=k_0}^{k_1} (2^{-j-1} \|\chi_{[0,t_j]} v | L^p\|)^p.$$

Together with (84), this implies

$$\|x_\varepsilon | L_v^p\|^p \leq \frac{2^{-p}(1+\varepsilon)^p}{1-2^{-p}} (2^{-k_0 p} \|\chi_{[0,t_{k_0}]} v | L^p\|)^p + \sum_{j=k_0}^{k_1} (2^{-j-1} \|\chi_{[t_j,t_{j+1}]} v | L^p\|)^p.$$

Next, using (83), we finally obtain

$$\|x_\varepsilon | L_v^p\| \leq \frac{(1+\varepsilon)}{(2^p-1)^{1/p}} \|x | L_v^p\|. \quad \square$$

Now, everything is ready for the proof of Theorem 1. We proceed with this.

Proof. By Lemma 2, the embedding

$$QL^p_{\bar{v}} \subseteq K(\downarrow) \cap L^p_v$$

follows from the boundedness of Q as an operator from $L^p_{\bar{v}}$ to L^p_v (that is how the weight \bar{v} was chosen). Furthermore, Q takes any nonnegative function to a monotone nonincreasing function. This proves (a).

We pass to the second statement of the theorem.

Fix a function $x \in K(\downarrow) \cap L^p_v$ with nonzero norm. Condition (11) implies the identity

$$(103) \quad \lim_{t \rightarrow \infty} x(t) = 0.$$

Since the norm in L^p_v is absolutely continuous, by (103) we see that there exists a strictly monotone continuous function $x_0 \in K(\downarrow) \cap L^p_v$ enjoying the conditions

$$(104) \quad x_0(t) \geq x(t) \quad (\forall t > 0);$$

$$(105) \quad \|x_0 | L^p_v\| \leq 2\|x | L^p_v\|;$$

$$(106) \quad \lim_{t \rightarrow \infty} x_0(t) = 0;$$

$$(107) \quad \lim_{t \rightarrow 0} x_0(t) = \infty.$$

Since the continuous function $x_0(t)$ is strictly monotone, by (106), (107) we deduce that for every $i \in \mathbb{Z}$ there exists a unique point $t_i \in (0, \infty)$ with $x_0(t_i) = 2^{-i}$. We define two new functions

$$y_{s0}(t) = \sum_{-\infty}^{\infty} \chi_{[t_i, t_{i+1})} 2^{-i}, \quad y_{s1}(t) = \sum_{-\infty}^{\infty} \chi_{[0, t_{i+1})} 2^{-i}.$$

Direct calculations show that for every $t > 0$ we have

$$(108) \quad y_{s1}(t) \equiv 2y_{s0}(t), \quad y_{s0}(t) \geq x_0(t) \geq \frac{1}{2}y_{s0}(t).$$

Consequently,

$$(109) \quad \|y_{s1} | L^p_v\| = 2\|y_{s0} | L^p_v\| \geq 2\|x_0 | L^p_v\| \geq \frac{1}{2}\|y_{s0} | L^p_v\|.$$

But from (105), (108), and the absolute continuity of the norm in L^p_v we deduce that

$$(110) \quad \begin{aligned} \lim_{i \rightarrow -\infty} \left\| \sum_{-\infty}^i \chi_{[0, t_{j+1})} 2^{-j} | L^p_v \right\| &= \lim_{i \rightarrow -\infty} 2 \left\| \sum_{-\infty}^i \chi_{[t_j, t_{j+1})} 2^{-j} | L^p_v \right\| \\ &= 2 \lim_{i \rightarrow -\infty} \|y_{s0} \chi_{[0, t_{j+1})} | L^p_v\| \leq 4 \lim_{i \rightarrow -\infty} \|x_0 \chi_{[0, t_{j+1})} | L^p_v\| = 0. \end{aligned}$$

Taking (110) into account, we apply Lemma 4 to y_{s1} . This yields a sequence $\{k_i\}_0^\infty$ such that

$$(111) \quad \left\| \sum_{k_0+1}^{\infty} 2^{-j} \chi_{[0, t_{j+1})} \mid L^p_v \right\| + \sum_{i=0}^{\infty} \left\| \sum_{k_{i+1}+1}^{k_i} 2^{-j} \chi_{[0, t_{j+1})} \mid L^p_v \right\| \leq 2 \|y_{s1} \mid L^p_v\|.$$

We define the functions

$$z_0 = \sum_{k_0+1}^{\infty} 2^{-j} \chi_{[0, t_{j+1})}; \quad z_{i+1} = \sum_{k_{i+1}+1}^{k_i} 2^{-j} \chi_{[0, t_{j+1})} \quad (i = 0, 1, 2, \dots).$$

Fixing $\varepsilon > 0$, we apply Lemma 6 to every function $\{z_i\}_0^\infty$. This will result in a collection of functions $\{z_{i\varepsilon}\}_0^\infty$ satisfying the conditions

$$(112) \quad Qz_{i\varepsilon}(t) \geq \frac{1}{16} z_i(t) \quad (\forall t > 0);$$

$$(113) \quad \|z_{i\varepsilon} \mid L^p_{\bar{v}}\| \leq \frac{1+\varepsilon}{(2^p-1)^{1/p}} \|z_i \mid L^p_v\|.$$

Put

$$x_\varepsilon = \sum_{i=0}^{\infty} z_{i\varepsilon}.$$

Then (104), (108), (111), and (112) show that for every $t > 0$ we have

$$Qx_\varepsilon(t) = Q\left(\sum_{i=0}^{\infty} z_{i\varepsilon}\right)(t) \geq \frac{1}{16} \sum_{i=0}^{\infty} z_i(t) = \frac{1}{16} y_{s1}(t) = \frac{1}{8} y_{s0}(t) \geq \frac{1}{8} x_0(t) \geq \frac{1}{8} x(t).$$

But from (105), (108), (109), and (113) we obtain

$$\begin{aligned} \|x_\varepsilon \mid L^p_{\bar{v}}\| &= \left\| \sum z_{i\varepsilon} \mid L^p_{\bar{v}} \right\| \leq \sum \|z_{i\varepsilon} \mid L^p_{\bar{v}}\| \leq \frac{1+\varepsilon}{(2^p-1)^{1/p}} \sum \|z_i \mid L^p_v\| \\ &\leq 2 \frac{1+\varepsilon}{(2^p-1)^{1/p}} \|y_{s1} \mid L^p_v\| \leq 8 \frac{1+\varepsilon}{(2^p-1)^{1/p}} \|x_0 \mid L^p_v\| \\ &\leq 16 \frac{1+\varepsilon}{(2^p-1)^{1/p}} \|x \mid L^p_v\|. \end{aligned} \quad \square$$

We pass to the proof of Theorem 7.

Proof. By Theorem 5, the cones $K(\downarrow) \cap L_v^\infty$ and $K(\downarrow) \cap L_{\bar{v}}^\infty$ coincide.

First, we verify (a). By Lemma 2, the operator $Q: L_u^\infty \rightarrow L_w^\infty$ is bounded if and only if

$$(114) \quad \sup_{t>0} w(t) \int_t^\infty \frac{ds}{u(s)} = C_8 < \infty.$$

In our case, (114) has the form

$$(115) \quad \sup_{t>0} \tilde{v}(t) \int_t^\infty \frac{ds}{\bar{v}(s)} = \sup_{t>0} \tilde{v}(t) \int_t^\infty \left(-\frac{d}{ds} \frac{1}{\bar{v}_{ac}}(s) \right) = \sup_{t>0} \tilde{v}(t) \frac{1}{\bar{v}_{ac}(t)} \leq C_8.$$

Now, we prove (b). Let $x \in K(\downarrow) \cap L_v^\infty$ and $c_0 = \sup_{t>0} x(t) \tilde{v}(t)$. Put $y_\varepsilon(t) = \frac{c_0}{\bar{v}(t)}$. Then $\|y_\varepsilon \mid L_{\bar{v}}^\infty\| = c_0$, and we have

$$\begin{aligned} Qy_\varepsilon(t) &= \int_t^\infty y_\varepsilon(\tau) d\tau = \int_t^\infty \frac{c_0}{\bar{v}(\tau)} d\tau \\ &= c_0 \int_t^\infty \left(-\frac{d}{ds} \frac{1}{\bar{v}_{ac}}(s) \right) ds = c_0 \frac{1}{\bar{v}_{ac}(t)}(t) \geq \frac{c_0}{c} x(t). \end{aligned} \quad \square$$

Now we pass to a lemma that yields a necessary and sufficient condition for the existence of an absolutely continuous function equivalent to a given weight function v .

We remark at once that, in the case of the space L_v^∞ , the value of the weight function at every point is essential. Therefore, we assume for definiteness in what follows that v is continuous from the left.

Lemma 7. *Fix $p = \infty$ and a weight function $v: [0, \infty) \rightarrow R_+$. Next, define a function \tilde{v} as in (34). Then the existence of an absolutely continuous function $\tilde{v}_{ac} \in K(\downarrow) \cap L_v^\infty$ such that*

$$(116) \quad \frac{1}{c} \tilde{v}_{ac}(t) \leq \tilde{v}(t) \leq c \tilde{v}_{ac}(t)$$

for some constant C and all $t \in R_+$ is equivalent to the inequality

$$(117) \quad \sup_{t \in R_+} \frac{\tilde{v}(t+0)}{\tilde{v}(t-0)} = d < \infty.$$

Proof. We verify the implication $(116) \Rightarrow (117)$. Since \tilde{v}_{ac} is absolutely continuous, by (116) we obtain

$$(118) \quad \frac{\tilde{v}(t+0)}{\tilde{v}(t-0)} \leq \frac{c \tilde{v}_{ac}(t+0)}{\tilde{v}(t-0)} \leq \frac{c^2 \tilde{v}_{ac}(t+0)}{\tilde{v}_{ac}(t-0)} = c^2,$$

and this proves the implication $(116) \Rightarrow (117)$.

Now, we verify the implication $(117) \Rightarrow (116)$. Since \tilde{v}_{ac} is monotone nondecreasing, the Radon–Nikodym decomposition yields the representation

$$(119) \quad \tilde{v}(t) = x_1(t) + x_2(t) + x_3(t),$$

where $x_1, x_2, x_3 \in K(\downarrow)$ are, respectively, absolutely continuous, singular, and jump functions. We construct an equivalent absolutely continuous function for each of them separately. The function x_1 is absolutely continuous itself, so nothing should be constructed for it. We have two possibilities for x_2 : a) this function is positive for all $t \in R_+$, b) there exists $\tau_0 \in R_+$ such that $x_2(\tau_0) = 0$ and $x_2(t) > 0$ for $t < \tau_0$. We consider a) first.

Put $c = x_2(1)$. For every $k \in Z$ we define $t_k \in R_+$ by $t_k = \sup\{t : x_2(t) \geq 2^k c\}$. If $0 \leq t_{k+1} < t_k < \infty$, then for $t \in [t_{k+1}, t_k]$ we define a function x_{2ac} by the formula $x_{2ac}(t) = x_2(t_k) + (t - t_k) \frac{x_2(t_{k+1}) - x_2(t_k)}{t_{k+1} - t_k}$. Then for all $t \in [t_{k+1}, t_k]$ we have

$$(120) \quad x_{2ac}(t_k) \leq x_2(t) \leq x_{2ac}(t_{k+1}) = 2x_{2ac}(t_k) \leq 2x_2(t).$$

If $0 < t_k < t_{k-1} = \infty$, then for $t \in [t_k, \infty)$ we define a function x_{2ac} by the formula $x_{2ac}(t) = x_2(t_k)$. Then for all $t \in [t_k, \infty)$ we have

$$(121) \quad x_2(t) \leq x_{2ac}(t_k) \leq 2x_2(t).$$

Now, if $x_2(0) < 2x_2(t_k)$, then for $t \in [0, t_k]$ we define x_{2ac} by the formula $x_{2ac}(t) = x_2(0)$. Then for $t \in [0, t_k]$ we again have (121).

Relations (120)–(121) show that in case a) we have constructed an absolutely continuous function x_{2ac} equivalent to x_2 with constant 2.

If condition b) is fulfilled, then for all $t \in [\tau_0, \infty)$ we define x_{2ac} by the formula $x_{2ac}(t) \equiv 0$. The subsequent construction is similar to case a) treated above.

Now, we show how to construct an equivalent absolutely continuous function in $K(\downarrow)$ for x_3 .

Again, there are two possibilities for x_3 : a) the function x_3 is positive for all $t \in R_+$, b) there exists $\tau_0 \in R_+$ such that $x_3(\tau_0) = 0$ and $x_3(t) > 0$ for $t < \tau_0$.

First, we consider case a). For every $k \in Z$, we define $t_k \in (\tau, \infty)$ by $t_k = \inf\{t : x_3(t) \leq (d+1)^k\}$. Again by continuity, $x_3(t_k) \leq (d+1)^k$ for all k .

Again, we construct a linear approximate. If $0 \leq t_{k+1} < t_k < \infty$, then for $t \in [t_{k+1}, t_k]$ we define a function x_{3ac} by the formula $x_{3ac}(t) = x_3(t_k) + (t - t_k) \frac{x_3(t_{k+1}) - x_3(t_k)}{t_{k+1} - t_k}$. Then, if $x_3(t_{k+1}) \leq (d+1)x_3(t_k)$, for all $t \in [t_{k+1}, t_k]$ we have

$$(122) \quad x_{3ac}(t_k) \leq x_3(t) \leq x_{3ac}(t_{k+1}) \leq (d+1)x_{3ac}(t_k) \leq (d+1)x_{3ac}(t).$$

But if $x_3(t_{k+1}) > (d+1)x_3(t_k)$, then for every $\delta > 0$ we have $x_3(t_k - \delta) > (d+1)^k$. Therefore, by (117) we obtain $x_3(t_{k+1}) \leq (d+1)^{k+1} \leq (d+1)x_3(t_k - \delta)$ and, consequently, $x_3(t_{k+1}) \leq d(d+1)x_3(t_k)$. Thus, for all $t \in [t_{k+1}, t_k]$ we have

$$(123) \quad x_{3ac}(t_k) \leq x_3(t) \leq x_{3ac}(t_{k+1}) \leq d(d+1)x_{3ac}(t_k) \leq d(d+1)x_{3ac}(t).$$

If $0 < t_k < t_{k-1} = \infty$, then for $t \in [t_k, \infty)$ we define x_{3ac} by the formula $x_{3ac}(t) = x_3(t_k)$.

Applying (117) once again, we see that

$$(124) \quad x_3(t) \leq x_{3ac}(t) \leq d(d+1)x_{3ac}(t)$$

for all $t \in [t_k, \infty)$. Relations (122)–(124) show that, in case a), we have constructed an absolutely continuous function x_{3ac} equivalent to x_3 with the constant $d(d+1)$.

If b) is fulfilled, then for all $t \in [t_0, \infty)$ we define x_{3ac} by $x_{3ac}(t) \equiv 0$. The subsequent arguments resemble those in case a). \square

In conclusion we discuss Lemma 8, required for the proof of Lemma 3.

Lemma 8. Fix $\theta \in (0, 1)$. In the spaces $L_{w_0}^1$ and $L_{w_1}^\infty$, consider the cones $K(\downarrow) \cap L_{w_0}^1$ and $K(\downarrow) \cap L_{w_1}^\infty$. Let

$$x \in (K(\downarrow) \cap L_{w_0}^1)^\theta (K(\downarrow) \cap L_{w_1}^\infty)^{1-\theta}$$

satisfy

$$(125) \quad \|x\| (K(\downarrow) \cap L_{w_0}^1)^\theta (K(\downarrow) \cap L_{w_1}^\infty)^{1-\theta} = 1.$$

Then there exists $x_0 \in K(\downarrow) \cap L_{w_0}^1$ with $\|x_0\| L_{w_0}^1 = 1$ such that

$$(126) \quad x(t) \leq x_0^\theta(t) \cdot \left(\frac{1}{\tilde{w}_1(t)} \right)^{1-\theta} (t)$$

for all $t \in R_+$.

Proof. Theorem 5 shows that the cones $K(\downarrow) \cap L_{w_1}^\infty$ and $K(\downarrow) \cap L_{\tilde{w}}^\infty$ coincide.

Suppose that (125) is fulfilled. This means that there exists a sequence of pairs of functions $x_{0n} \in K(\downarrow) \cap L_{w_0}^1$ with $\|x_{0n}\| L_{w_0}^1 = 1$ and $x_{1n} \in K(\downarrow) \cap L_{\tilde{w}}^\infty$ with $\|x_{1n}\| L_{\tilde{w}}^\infty = 1$ such that for all $t \in R_+$ we have

$$(127) \quad x(t) \leq \left(1 + \frac{1}{n}\right) \cdot x_{0n}^\theta(t) x_{1n}^{1-\theta}(t).$$

Since every $x \in K(\downarrow) \cap L_{w_1}^\infty$ with $\|x\| L_{w_1}^\infty = 1$ satisfies $x(t) \leq \frac{1}{\tilde{w}(t)}$ for all $t \in R_+$, we can rewrite (127) in the form

$$(128) \quad x(t) \leq \left[\left(1 + \frac{1}{n}\right)^{1/\theta} \cdot x_{0n}(t) \right]^\theta \cdot \left(\frac{1}{\tilde{w}_1(t)} \right)^{1-\theta}.$$

We define a function y at all points $t \in R_+$ by the formula $y(t) = \inf_n \left(1 + \frac{1}{n}\right)^{1/\theta} \cdot x_{0n}(t)$. Then $y \in K(\downarrow) \cap L_{w_0}^1$, and the pointwise inequality $y(t) \leq \left(1 + \frac{1}{n}\right)^{1/\theta} \cdot x_{0n}(t)$ shows that $\|y\| L_{w_0}^1 \leq \left(1 + \frac{1}{n}\right)^{1/\theta}$ for every $n \in N$. This means that $\|y\| L_{w_0}^1 \leq 1$. In (128), we pass to the infimum over n for every $t \in R_+$, obtaining the inequality

$$(129) \quad x(t) \leq y(t)^\theta \cdot \left(\frac{1}{\tilde{w}_1(t)} \right)^{1-\theta}$$

valid for all $t \in R_+$. Now, if we suppose that $\|y\|_{L_{w_0}^1} = q < 1$, then, putting $q_0 = q^\theta$, we see that for all $t \in R_+$ we have

$$x(t) \leq \left(\frac{y(t)}{q}\right)^\theta \cdot \left(\frac{q_0}{\tilde{w}_1(t)}\right)^{1-\theta};$$

moreover, $\frac{q_0}{q}y \in K(\downarrow) \cap L_{w_0}^1$ with $\|\frac{q_0}{q}y\|_{L_{w_0}^1} = q_0$, and also $q_0 \frac{1}{\tilde{w}_1} \in K(\downarrow) \cap L_{\tilde{w}_1}^\infty$ with $\|q_0 \frac{1}{\tilde{w}_1}\|_{L_{\tilde{w}_1}^\infty} = q_0$, which contradicts (125). \square

REFERENCES

- [1] M. A. Ariño and B. Muckenhoupt, *Maximal functions on classical Lorentz spaces and Hardy's inequality with weights for nonincreasing functions*, Trans. Amer. Math. Soc. **320** (1990), no. 2, 727–735. MR989570
- [2] E. I. Bereznoi, *Interpolation of positive operators in the spaces $\phi(X_0, X_1)$* (Russian), Qualitative and approximate methods for the investigation of operator equations (Russian), Yaroslav. Gos. Univ., Yaroslavl', 1981, pp. 3–12, 160. MR881222
- [3] E. I. Bereznoi, *Two-weight estimates for a class of integral operators*, Tr. Mat. Inst. Steklov. **201** (1992), 14–25; English transl., Proc. Steklov Inst. Math. **201** (1994), no. 2, 11–20. MR1291988 (95i:47057)
- [4] E. I. Bereznoi, *Sharp estimates for operators on cones in ideal spaces*, Tr. Mat. Inst. Steklov. **204** (1993), 3–34; English transl., Proc. Steklov Inst. Math. **204** (1994), no. 3, 1–25. MR1320016 (96a:46052)
- [5] E. I. Bereznoi, *Theorems on the representation of spaces and Schur's lemma*, Dokl. Akad. Nauk **344** (1995), no. 6, 727–729; English transl., Dokl. Math. **52** (1995), 252–254. MR1364314 (97b:47030)
- [6] E. I. Bereznoi, *Two-weighted estimations for the Hardy-Littlewood maximal function in ideal Banach spaces*, Proc. Amer. Math. Soc. **127** (1999), no. 1, 79–87. MR1622773
- [7] E. I. Bereznoi, *Schur test for the Hardy operator*, Eurasian Math. J. **4** (2013), no. 4, 17–30. MR3382900
- [8] E. I. Bereznoi and L. Maligranda, *Extrapolation of operators and the limits of applicability of Schur's test*, Dokl. Akad. Nauk **393** (2003), no. 5, 583–586; English transl., Dokl. Math. **68** (2003), no. 3, 392–395. MR2089642 (2005f:47114)
- [9] E. I. Bereznoi and L. Maligranda, *Representation of Banach ideal spaces and factorization of operators*, Canad. J. Math. **57** (2005), no. 5, 897–940. MR2164589
- [10] E. I. Bereznoi and L. Maligranda, *Representability of some cones in weighted Lebesgue spaces and the extrapolation of operators on cones*, Dokl. Akad. Nauk **406** (2006), no. 4, 439–442; English transl., Dokl. Math. **73** (2006), no. 1, 59–62. MR2347324 (2008i:46026)
- [11] J. Bergh and J. Löfström, *Interpolation spaces. An introduction*, Springer-Verlag, Berlin-New York, 1976. Grundlehren der Mathematischen Wissenschaften, No. 223. MR0482275
- [12] Yu. A. Brudnyi and N. Ya. Krugljak, *Interpolation functors and interpolation spaces. Vol. I*, North-Holland Mathematical Library, vol. 47, North-Holland Publishing Co., Amsterdam, 1991. Translated from the Russian by Natalie Wadhwa; With a preface by Jaak Peetre. MR1107298
- [13] A.-P. Calderón, *Intermediate spaces and interpolation, the complex method*, Studia Math. **24** (1964), 113–190. MR0167830
- [14] M. J. Carro and J. Soria, *Weighted Lorentz spaces and the Hardy operator*, J. Funct. Anal. **112** (1993), no. 2, 480–494. MR1213148
- [15] M. Christ, *Weighted norm inequalities and Schur's lemma*, Studia Math. **78** (1984), no. 3, 309–319. MR782668
- [16] J. García-Cuerva and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North-Holland Mathematics Studies, vol. 116, North-Holland Publishing Co., Amsterdam, 1985. Notas de Matemática [Mathematical Notes], 104. MR807149
- [17] A. Gogatishvili and V. D. Stepanov, *Reduction theorems for weighted integral inequalities on the cone of monotone functions* (Russian, with Russian summary), Uspekhi Mat. Nauk **68** (2013), no. 4 (412), 3–68; English transl., Russian Math. Surveys **68** (2013), no. 4, 597–664. MR3154814
- [18] M. L. Gol'dman, *Order-sharp estimates for Hardy-type operators on cones of quasimonotone functions*, Eurasian Math. J. **2** (2011), no. 3, 143–146. MR2910846
- [19] M. L. Gol'dman, H. P. Heinig, and V. D. Stepanov, *On the principle of duality in Lorentz spaces*, Canad. J. Math. **48** (1996), no. 5, 959–979. MR1414066
- [20] H. Heinig and L. Maligranda, *Weighted inequalities for monotone and concave functions*, Studia Math. **116** (1995), no. 2, 133–165. MR1354136

- [21] H. P. Heinig and V. D. Stepanov, *Weighted Hardy inequalities for increasing functions*, Canad. J. Math. **45** (1993), no. 1, 104–116. MR1200323
- [22] A. Kamińska and L. Maligranda, *Order convexity and concavity of Lorentz spaces $\Lambda_{p,w}$, $0 < p < \infty$* , Studia Math. **160** (2004), no. 3, 267–286. MR2033403
- [23] A. Kamińska and M. Mastyło, *Abstract duality Sawyer formula and its applications*, Monatsh. Math. **151** (2007), no. 3, 223–245. MR2329084
- [24] L. V. Kantorovich and G. P. Akilov, *Functional Analysis*, 2nd ed., Nauka, Moscow, 1977; English transl., Pergamon Press, Oxford, 1982. MR0664597 (83h:46002)
- [25] V. B. Korotkov, *Integral operators*, Nauka, Novosibirsk, 1983. (Russian). MR0721475 (85g:47071)
- [26] S. Krein, Yu. I. Petunin, and E. M. Semenov, *Interpolation of linear operators*, Nauka, Moscow, 1978; English transl., Transl. Math. Monogr., vol. 54, Amer. Math. Soc., Providence, RI, 1982. MR0649411 (84j:46103)
- [27] A. Kufner, L. Maligranda, and L.-E. Persson, *The Hardy inequality. About its history and some related results*, Vydavatelský Servis, Plzeň, 2007. MR2351524
- [28] A. Kufner and L.-E. Persson, *Weighted inequalities of Hardy type*, World Scientific Publishing Co., Inc., River Edge, NJ, 2003. MR1982932
- [29] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces. II. Function spaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas], vol. 97, Springer-Verlag, Berlin-New York, 1979. MR540367
- [30] L. Maligranda, *Calderón–Lozanovskii spaces and interpolation of operators*, Semesterbericht Funktionalanalysis **8** (1985), 83–92.
- [31] L. Maligranda, *Orlicz spaces and interpolation*, Semin. Math., vol. 5, Univ. Campinas, Campinas, 1989. MR2264389 (2007e:46025)
- [32] L. Maligranda, *Weighted estimates of integral operators for decreasing functions*, Boundary value problems, special functions and fractional calculus (Minsk, 1996), Belarus. Gos. Univ., Minsk, 1996, pp. 226–236. MR1428945
- [33] L. Maligranda, *Weighted inequalities for monotone functions*, Collect. Math. **48** (1997), no. 4-6, 687–700. MR1602636
- [34] L. Maligranda, *Weighted inequalities for quasi-monotone functions*, J. London Math. Soc. (2) **57** (1998), no. 2, 363–370. MR1644209
- [35] V. G. Maz'ya, *Sobolev Spaces*, Leningrad. Univ., Leningrad, 1985; English transl., Springer Ser. Soviet Math., Springer-Verlag, Berlin, 1985. MR0807364 (87g:46055)
- [36] E. Sawyer, *Boundedness of classical operators on classical Lorentz spaces*, Studia Math. **96** (1990), no. 2, 145–158. MR1052631
- [37] V. A. Shestakov, *Transformations of Banach ideal spaces and interpolation of linear operators* (Russian, with English summary), Bull. Acad. Polon. Sci. Sér. Sci. Math. **29** (1981), no. 11-12, 569–577 (1982). MR654216
- [38] G. Sinnamon, *Transferring monotonicity in weighted norm inequalities*, Collect. Math. **54** (2003), no. 2, 181–216. MR1995140
- [39] G. Sinnamon, *Hardy's inequality and monotonicity*, Function Spaces and Nonlinear Analysis, Math. Inst. Acad. Sci. Czech Republic, Prague, 2005, pp. 292–310.
- [40] V. D. Stepanov, *Integral operators on the cone of monotone functions*, J. London Math. Soc. (2) **48** (1993), no. 3, 465–487. MR1241782
- [41] V. D. Stepanov, *The weighted Hardy's inequality for nonincreasing functions*, Trans. Amer. Math. Soc. **338** (1993), no. 1, 173–186. MR1097171
- [42] P. Szeptycki, *Notes on integral transformations*, Dissertationes Math. (Rozprawy Mat.) **231** (1984), 48 pp. MR763933

P. G. DEMIDOV YAROSLAVL, STATE UNIVERSITY, UL. SOVETSKAYA 14, 150000 YAROSLAVL, RUSSIA
 Email address: ber@uniyar.ac.ru

LULEÅ UNIVERSITY OF TECHNOLOGY, S-971 87, LULEÅ, SWEDEN
 Email address: lech.maligranda@ltu.se

Received 15/MAY/2016

Translated by S. V. KISLYAKOV