
Linköping University Post Print

Computerized lifestyle intervention in routine primary health care: Evaluation of usage on provider and responder levels



N.B.: When citing this work, cite the original article.

Original Publication:

Siw Carlfjord, Per Nilsen, A Andersson, Kjell Johansson and Preben Bendtsen, Computerized lifestyle intervention in routine primary health care: Evaluation of usage on provider and responder levels, 2009, PATIENT EDUCATION AND COUNSELING, (75), 2, 238-243.

http://dx.doi.org/10.1016/j.pec.2008.10.004 Copyright: Elsevier Science B.V., Amsterdam. http://www.elsevier.com/

Postprint available at: Linköping University Electronic Press http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-18143

Computerized lifestyle intervention in routine primary care: Evaluation of usage on provider and responder levels

S. Carlfjord^a,*, P. Nilsen^a, M. Leijon^{a,b}, A. Andersson^{a,c}, K. Johansson^a, P. Bendtsen^a

^aDepartment of Medical and Health Sciences, Linköping University, Linköping, Sweden

^bCentre for Public Health and Sciences, County Council of Östergötland, Linköping, Sweden

^cCounty Council of Östergötland, R&D Department of Local Health Care, Linköping, Sweden

*Corresponding author at: Department of Medical and Health Sciences, Division of

Community Medicine, Linköping University, SE-581 83 Linköping, Sweden. Tel.:

+46(0)13227132 fax: +46(0)13149403. E-mail address: siw.carlfjord@ihs.liu.se (S.

Carlfjord).

Abstract

Objective

The aim of this study was to evaluate the use of a computerized concept for lifestyle intervention in routine primary health care (PHC).

Methods

Nine PHC units were equipped with computers providing a lifestyle test and tailored printed advice. Patients were referred by staff, and performed the test anonymously. Data were collected over a period of one year.

Results

An average of 5.7% of the individuals visiting PHC during the study period performed the test. There were great differences between the units in the number of tests performed and in the proportion of patients referred. One-fifth of the respondents scored for hazardous alcohol consumption, and one-fourth reported low levels of physical activity. A majority of respondents found the test easy to perform, and a majority of those referred to the test found referral positive.

Conclusion

The computerized test can be used for screening and intervention regarding lifestyle behaviour in primary care. Responders are positive to the test and to referral.

Practice implications

A more widespread implementation of computerized lifestyle tests could be a beneficial complement to face-to-face interventions in primary care.

Keywords: Alcohol; Computer-based; Lifestyle intervention; Physical activity; Primary health care

1. Introduction

Lifestyle behaviours such as alcohol consumption, smoking, diet and physical activity have been shown to have a great impact on health [1–3]. Individuals can make important contributions to their own health by adopting some health-related behaviours and avoiding others [4–6]. In many countries, lifestyle related diseases are the leading causes of death [7–9]. There is an increasing awareness among health care authorities that unhealthy lifestyle behaviours need to be addressed in routine health care. However, most health care systems focus on the management of acute illness and chronic health conditions. Health care providers face substantial barriers to providing preventive services aimed at achieving health-related behaviour change, as they often lack the time, knowledge, and skills [10–11].

The use of computer technology has been suggested as a way to overcome many of the barriers to integrating health behaviour change interventions into routine health care.

Computer-based screening and advice, office-based or web-based, for various health-related behaviours has been developed during recent decades and there is a growing body of evidence supporting its effectiveness [12-15]. In a study by Kypri et al. [16], computerized alcohol interventions performed as well as practitioner-delivered brief interventions.

Moreover, computerized interventions concerning alcohol or physical activity have been favourably evaluated in terms of feasibility, provider acceptability and patient willingness to participate in various settings, including emergency departments, primary care, and schools [17–22].

Research has demonstrated that computer-assisted health behaviour advice may have several advantages compared to conventional face-to-face counselling. The use of computers has been found to decrease the effect of social desirability and increase the amount of information disclosed [23–24]. Findings suggest that patients prefer to reveal information of a

personal and potentially embarrassing nature to a computer rather than a person [25–27]. The use of computer-based screening and advice can improve the effectiveness of behavioural counselling through improved consistency of counselling and closer matching of intervention to patient characteristics and recommended guidelines. The number of staff needed to deliver counselling and the associated costs for personnel training can be reduced when advice is delivered by a computer [28].

Despite expanding documentation on the effectiveness and feasibility of delivering computer-based lifestyle interventions in health care, there is a paucity of research that evaluates implementation of such interventions in primary health care (PHC). The aim of this study was to evaluate the use of a computerized concept for lifestyle testing and tailored advice implemented in routine PHC, in terms of provider usage, responder characteristics and responder attitudes to the concept. To the best of our knowledge, this is the first attempt to evaluate a computerized lifestyle intervention concept in a routine PHC setting.

2. Methods

2.1. Setting

The study was conducted in Östergötland, Sweden, a county with approximately 420,000 inhabitants, considered representative of the Swedish population in terms of age distribution, employment rates and economy. Nine PHC units, from different areas, urban and rural areas, were recruited to the study, which started in autumn 2006. The units also varied in the number of general practitioners (GPs), nurses and other staff members employed. The size of the PHC units, measured by number of listed patients aged18 years or more, ranged from 4200 to 16,500 (average 9500). The study was carried out as a development project in routine PHC, in cooperation with the health care authorities in Östergötland County. In Sweden, health care is

publicly funded, i.e. residents are insured by the state and health care services are funded through a taxation scheme of county councils.

2.2. The computerized concept

The computer-based lifestyle intervention concept was developed by the Lifestyle Intervention Research group at Linköping University. It was based on previous experiences from student health care and emergency department settings, as reported in Karlsson and Bendtsen [17], Bendtsen et al. [18] and Karlsson et al. [29].

The lifestyle test included questions on the following topics: alcohol consumption; physical activity; motivation to change; and attitudes to performing the test. The questions on alcohol consumption were beverage-specific and evaluated weekly consumption on a day-by-day basis and frequency of heavy episodic drinking (HED), i.e. intake of a large volume of alcohol on any one occasion. If the respondent reported no alcohol consumption during the last 3 months, the subsequent alcohol questions were omitted. If a patient reported they had been referred to the test, they were also asked about which staff category made the referral and about their attitude to being referred. Respondents who completed the test received a printed sheet with tailored written advice based on their answers and categorization of the level of alcohol consumption as "hazardous", "increased risk" or "low risk" and the level of physical activity as "inactive", "insufficiently active" or "active".

Alcohol consumption was measured by number of standard drinks (12 g alcohol) per week, and frequency of HED. Hazardous consumption was defined for a woman as 10 or more standard drinks per week and/or 4 standard drinks per occasion (HED) once a week or more frequently, and for a man 15 or more standard drinks per week and/or 5 standard drinks per occasion (HED) once a week or more frequently. Physical activity questions were based on recommendations from the Centers for Disease Control and Prevention (CDC) and the

American College of Sports Medicine in 1995 [30]. The questions measured number of days per week with moderate-intensity aerobic (endurance) physical activity for a minimum of 30 min (rendered 1 point/day), and number of days per week with vigorous-intensity aerobic physical activity (rendered 2 points/day). To be considered physically active, 5 points had to be obtained. The questions concerning motivation to change alcohol consumption were influenced by the Stages of Change model and categorized the respondents who had no intention of changing as pre-contemplators, those considering change as contemplators, and those who had made a decision to change their habits as being in preparation phase [31].

2.3. Implementation activities

Managers and health coordinators at the nine PHC units were invited to an information session with the research team and agreed to participate in the project. All staff members were then invited to an information session at their own PHC unit, where the test was demonstrated and they were educated about how it should be used. Since it was not deemed realistic to expect all patients visiting the health care centre to be referred to the test the staff were told to decide which patients should be the target group for the test, simply in order to avoid forgetting to use the test. Thus, at each unit staff were allowed to choose the patient groups that should be included in the testing routine, as well as the staff categories that should refer patients to the test. Hence, the patient groups and the referring staff categories varied between the units. All were told to refer patients who had their blood pressure tested. Four units added one group (aged 20–30 or "not severe mental disorders"), one unit added two more groups (aged 20–30 and aged 45–65), and one unit selected five groups in addition to "blood pressure" (blood lipid testing, aged 20–30, "not severe mental disorders", smokers, "patients on sick leave >28 days). The remaining three units did not add any groups. An agreement stating which patient groups and staff categories for referral were chosen was signed by the

manager at each unit. At least one person at each participating unit was appointed as a contact person between the PHC unit and the research team.

The nine PHC units were provided with computers, monitors and printers integrated in an IT kiosk specially designed for the project. PHC units with less than 11,000 listed patients aged 18 or older received one computer kit, larger units (three of the nine included) received two computer kits. The computers were equipped with the lifestyle test and the IT kiosks were placed in waiting rooms, corridors or in a separate room, depending on local conditions.

After an initial period of learning how to use the test, the computers were connected to the local health care computer network in Östergötland County, which made it possible to store data centrally, and give regular feedback to the participating units. From April 2007 each PHC unit was provided with weekly statistical feedback from the test, including the number of tests performed, proportion of respondents who stated they had been referred to the test by a staff member, and by which staff category, and proportion of respondents with hazardous alcohol consumption and/or low physical activity. Members from the research team visited the participating units now and then during the implementation period, to check out the computers were active, but also to participate in staff meetings. Reminder posters were distributed to all units to help staff remember to refer their patients to the test.

The objective was that staff members should actively refer the patients to the test. However, the patients were also free to spontaneously initiate tests. GPs, nurses or other staff members invited patients to perform the test after the normal consultation. Patients below 18 years of age, not able to understand Swedish, or too ill were excluded from referral. Staff members were asked to report in the medical records when a patient had been advised to perform the test. The test was performed anonymously, and the respondents themselves decided whether to discuss the test results with any member of the PHC staff. It was assumed

that the majority of the respondents would not ask for additional help to change their healthrelated behaviours, but to take responsibility for their own health. However, if a patient asked for extended counselling the PHC unit had agreed to provide some consultation.

2.4. Data collection and analysis

The data were analysed after one year of data collection from April 2007 to March 2008. Data concerning the number of listed patients at each unit and the number of individuals who visited the PHC during the period were collected from county council files. All other data were obtained from the test results. Statistical analyses were performed using the computer-based analysis program SPSS (Statistical Package for the Social Sciences) version 15.0. Analyses were performed using the chi-squared test or the Kruskal–Wallis test. Correlations were analyzed using Pearson's correlation coefficient. Statistical significance was defined as a p-value ≤ 0.05 .

3. Results

During the one-year study period, 5202 tests were initiated, of which 3065 tests (59%) were completed. Among the completed tests, 38 respondents stated that they had consumed >100 standard drinks per week. These were regarded outliers and were not included in the data analysis. No outliers in terms of physical activity were found. The remaining 3027 tests were included in the further analyses.

3.1. Usage on a provider level

Table 1 shows the number of tests performed, either by patients who were referred by staff or spontaneously without referral. The decrease in July occurred during the most common vacation period in Sweden. The average proportion of tests in relation to the total number of individuals who attended the PHC units during the study period was 5.7%.

Table 1 Number of respondents who performed the test month by month during the one-year study period (starting April 2007), referred or spontaneously, at the nine participating PHC units

	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Total
Number of tests performed	276	345	244	128	207	287	345	320	224	233	234	184	3027
Referred ^a	116	151	101	24	71	138	184	151	95	107	98	84	1320
Not referred ^b	160	194	43	04	136	149	161	169	129	126	136	100	1707

^aRespondent stated that he or she was referred to the test by a staff member.

The characteristics of the nine PHC units in terms of the size of the units, selected referral groups, number of tests and referrals, are shown in Table 2. There were considerable differences not only in the number of tests performed, but also in the proportion of respondents who were actively referred by the staff to perform the test. Forty-four percent (range 11–87% per unit) of the completed tests were performed by individuals who stated that they had been referred by staff. These represent 2.5% of the individuals who visited their PHC unit during the study period (range 0.7–8.3% per unit). The unit that selected the highest number of referral groups had one of the lowest proportions of referred patients.

3.2. Responder characteristics

Slightly more than half of the tests (52%) were completed by women. One-fifth of the respondents scored for hazardous alcohol consumption (range 16–23% per unit) and one-fourth (range 21–30% per unit) of the respondents had low levels of physical activity (insufficiently active 13%, inactive 14%). Men had higher rates (p<0.001) of hazardous alcohol consumption (28%) than women (14%), and lower rates (p<0.05) of being physically active (72%) than women (75%).

^bRespondent stated that he or she was not referred to the test by a staff member.

Table 2 Size of PHC unit, referral groups, computer localization, number of tests and active referrals to the test

Unit	Listed patients ≥18	Individuals aged ≥18 who attended the PHC ^a	Number of groups selected ^b	Number of completed tests	Number of respondents actively referred to the test	Proportion completed tests/individuals attending the PHC (%) ^a	Proportion referred/total number of completed tests (%)	Proportion referred/individuals attending the PHC (%) ^a
A ^c	4209	3277	2	232	128	7.1	55	3.9
$B^{d} \\$	5612	3592	3	281	246	7.8	87	6.8
C^c	6179	4063	2	453	339	11.1	75	8.3
D^{e}	7598	4738	2	170	82	3.6	48	1.7
E ^c	9127	4907	2	291	79	5.9	27	1.6
F^c	10766	6744	1	260	58	3.9	22	0.9
G^c	11223	6049	1	282	148	4.7	52	2.4
H ^c	14421	8798	1	368	167	4.2	45	1.9
I ^e	16509	10720	6	690	73	6.4	11	0.7
Total	85644	52888	n.r.	3027	1320			
Average						5.7	44	2.5

^aDuring the one-year study period, starting March 2007.

Sixty percent of the hazardous alcohol consumers were, according to the Stages of Change model, identified as pre-contemplators, 24% as contemplators, and 16% as being in the preparation phase for change [31]. There were no differences according to gender.

Three-fourths of the respondents stated that they intended to increase their physical activity level and one-fourth did not express such an intention (only two choices were available).

Those already physically active were significantly more interested in increasing their current physical activity than those who were categorized as insufficiently active or inactive

^bEach unit selected groups of patients who should be referred to the test.

^cComputer placed in corridor.

^dComputer placed in separate room.

^eComputer placed in waiting room.

(p<0.001), the proportions were 56% among those insufficiently active or inactive, and 82% among the physically active. No gender differences were found.

Those who had been referred to perform the test (44%) were significantly older than those performing the test spontaneously (p<0.001). Almost half of the referred participants (43%) were aged 60 years or older, 21% were aged 51–60, 14% were aged 41–50, 10% were aged 31–40, 6% were aged 21–30 and 5% were from the youngest age group, 18–20 years old. With regard to levels of hazardous consumption or physical activity, no differences were found between respondents who were referred by staff and those performing the test spontaneously. The proportion of spontaneous tests was positively associated with the proportion of interrupted tests at each unit (r=0.8, p<0.05).

3.3. Responder attitudes to the test

The vast majority (88%) of the respondents who completed the test found it easy or very easy to perform, with no correlations with gender or age. Referred patients found the test significantly easier to perform than those who performed the test spontaneously (p<0.001). Of the referred patients 90% found it easy, and 3% had difficulties; 5% of the non-referred reported difficulties and 87% found the test easy to perform. Among participants with hazardous alcohol consumption, 7% reported difficulties; 4% of the non-hazardous consumers perceived difficulties in performing the test. This difference was not significant (p=0.056).

The majority of patients (84%) who were referred to the test found referral positive, whereas 3% found it negative to be referred. Men were less positive to the referral than the women, with 4% of men, and 2% of women reporting that they found it negative to be referred (p<0.01). Respondents with hazardous alcohol consumption (p<0.001) or low physical activity levels (p<0.05) found it significantly less positive to be referred. Among the hazardous alcohol consumers, 6% were negative, and among others 2%. Among the inactive

or insufficiently physically active respondents, 4% were negative to the referral; 2% of the physically active respondents had a negative attitude to being referred to the test.

4. Discussion and conclusion

4.1. Discussion

This study was conducted in order to investigate the usage of a computer-based concept for assessment of and tailored advice concerning alcohol consumption and physical activity, when implemented in routine PHC. The results showed considerable variation between the participating PHC units with regard to the proportion of patients who performed the test and the proportion of respondents who reported having been referred to the test, despite similar implementation efforts. Referral rates were likely affected by many factors not investigated in this study, but the placement of the computer, the size of the unit and the number of selected referral groups appears to have impacted on the proportion of patients being referred.

Interestingly, the unit that selected the highest number of patient groups to refer actually had the lowest proportion of referred patients of all units. This could suggest that a high number of selected groups leads to confusion among staff members, and thus obstructs the referral process. In terms of hazardous behaviour regarding alcohol consumption or physical activity, there were no differences between referred patients and those who performed the test spontaneously.

An average of 5.7% of all individuals attending the nine PHC units during the one-year study period completed the test. This proportion could probably have been higher if more effort had been made to encourage the staff to refer patients to the test. However, the purpose was to examine the implementation of a new working method under real-world conditions, i.e. studying implementation effectiveness rather than efficacy. The implementation decisions were a top-down process, with managers deciding whether to participate in the project. It is

possible that a higher degree of staff involvement in the decision phase would have resulted in higher referral rates, and thus, in a higher proportion of individuals performing the test. In a study conducted in Virginia, USA, where primary care clinicians referred their patients to a web site that provided information for behavioural change, 4% of patients who had attended the participating family care practices visited the web site [32]. Implementation success has been shown to depend on the characteristics of the innovation, adopter characteristics and the context in which the implementation occurs [33]. In our study the innovation itself did not differ between the units, but adopter characteristics and the context varied to a certain degree.

One aim of the study was to investigate responder attitudes to the computerized test. They were found to be mainly positive. The respondents stated that they found the test easy to use and those referred to the test were predominantly positive to the referral, which implicates a feasible concept at the responder level. However, those most negative to being referred to the test probably did not answer the test at all, or initiated it but did not complete it. It is also possible that respondents who had difficulties in performing the test were more likely to leave it unfinished than those who found it easy.

Attitudes to the referral differed between men and women, with men expressing less positive attitudes to being referred. Past research has shown that men are more likely to change their alcohol consumption after brief interventions than women [34]. However, men do have higher alcohol consumption, and consumption levels were associated with a more negative attitude to referral to the test. A similar pattern was seen with regard to physical activity; those respondents who had low levels of physical activity were more negative to being referred than those who were physically active. It is possible that awareness of having a less healthy lifestyle predicts a more reluctant attitude to being referred to the test.

The proportion of respondents reporting hazardous alcohol consumption in our study was higher than in former studies in Swedish settings although the measurement methodology differed somewhat [35,36]. The levels of physical activity found coincided quite well with former studies in Sweden [37]. Since the target group consisted of primary care patients, who are older than the population in general, low alcohol consumption levels and low levels of physical activity could be expected. The alcohol consumption levels reported could possibly be a result of more honest self-reports from respondents using the computer, instead of a face-to-face counselling session. Previous research has demonstrated that the use of computerized self-administered assessment tools results in higher reporting of socially undesirable behaviours [26].

A considerable proportion of the respondents stated that they were prepared to change their habits. According to the test results, more than one-third of the hazardous alcohol consumers were thinking about changing their alcohol consumption habits, and more than half of those reporting low levels of physical activity were determined to increase this. This information about the patients could possibly help primary care staff in their efforts to guide and support their patients to a lifestyle change. In terms of physical activity, one way of doing this could be the method of "physical activity on prescription", which has been evaluated and found feasible in various settings [38,39]. According to Pinto et al. [5], addressing lifestyle factors should be a "gold standard" in primary care. When doctors, nurses, or other staff groups refer their patients to the lifestyle test, it is a way of emphasizing to the patient the medical staff's concerns about the influence of lifestyle factors on health. There is also a potential learning effect for the staff, since discussions about the feedback concerning hazardous lifestyle among patients could lead to an increased awareness of the importance of supporting the patients to choose a healthy lifestyle, and in the next step to new routines in everyday care, i.e. a double-loop learning process [40].

4.2. Study limitations

One limitation of this study is that there is no information about those patients who were referred to the test but did not do it, or did not complete it. Another limitation is that only a small number of implementation aspects were considered.

4.3. Conclusions

The study shows that the computerized test can be used for screening and intervention regarding lifestyle behaviour in a primary care setting, even though a fairly low proportion of the patients were referred to perform the test. The responder characteristics revealed high levels of hazardous behaviour regarding alcohol consumption. Responder attitudes to the test and to the referral were generally positive, thus the concept could be considered feasible at responder level.

4.4. Practice implications

The computerized concept offers a means of providing lifestyle intervention in PHC. A more widespread implementation of computerized lifestyle tests could be a beneficial complement to face-to-face interventions.

Acknowledgements

The study was supported by the county council of Östergötland, Sweden. The authors are grateful to Mrs Lena Lindhe-Söderlund for support in developing and designing the computerized lifestyle test, to Mr Mikael Åkeborg for computer programming and to Mrs Marika Holmqvist for statistical support.

References

- [1] WHO. The world health report 2002 Reducing risks, promoting healthy life. Geneva: World Health Organization; 2002. Available at http://www.who.int
- [2] Tones K, Green J. Health promotion. London: Sage Publications; 2004.
- [3] Brønnum-Hansen H, Juel K, Davidsen M, Sørensen J. Impact of selected risk factors on expected lifetime without long-standing, limiting illness in Denmark. Prev Med 2007;45:49–53.
- [4] Rosal MP, Ockene JK, Luckmann R, Zapka J, Goins KV, Saperia G, Mason T, Donnelly G. Coronary heart disease multiple risk factor reduction: providers' perspectives. Am J Prev Med 2004;27(2 Suppl 1):54–60.
- [5] Pinto BM, Rabin C, Farrell N. Lifestyle and coronary heart disease prevention. Prim Care 2005;32:947–61.
- [6] Conner M, Norman P. Predicting health behaviours. 2nd ed. Buckingham, UK: Open University Press; 2005.
- [7] Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. JAMA 2004;291:1238–45.
- [8] Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Correction: actual causes of death in the United States, 2000. JAMA 2005;293:293–4.
- [9] Alfredsson J. The development and distribution of public health. Health in Sweden: The National Public Health Report 2005. Scand J Public Health 2006;34: 27–42.

- [10] Johansson K, Bendtsen P, Åkerlind I. Early intervention for problem drinkers: readiness to participate among general practitioners and nurses in Swedish primary health care. Alcohol Alcohol 2002;37:38–42.
- [11] Stange K, Woolf S, Gjeltema K. One minute for prevention. The power of leveraging to fulfill the promise of health behavior counseling. Am J Prev Med 2002;22:320–3.
- [12] Kreuter MW, Strecher VJ. Do tailored behavior change messages enhance the effectiveness of health risk appraisal? Results from a randomized trial. Health Educ Res 1996;11:97–105.
- [13] Brug J, Campbell M, van Assema P. The application and impact of computer-generated personalized nutrition education: a review of the literature. Patient Educ Couns 1999;36:145–56.
- [14] Kypri K, Stephenson S, Langley J, Cashell-Smith M, Saunders J, Russell D. Computerised screening for hazardous drinking in primary care. N Z Med J 2005;118:U1703.
- [15] Martin K, Jarvis LA, Beale B. Healthy lifestyle check. A computerized health screening program. Comput Nurs 1997;15:77–81.
- [16] Kypri K, Saunders J, Williams SM, McGee RO, Langley JD, Cashell-Smith ML, Gallagher SJ. Web-based screening and brief intervention for hazardous drinking: a double-blind randomized controlled trial. Addiction 2004;99:1410–17.
- [17] Karlsson A, Bendtsen P. Acceptability of a computerized alcohol screening and advice routine in an emergency department setting a patient perspective. Addict Behav 2005;30:767–76.

- [18] Bendtsen P, Holmqvist M, Johansson, K. Implementation of computerized alcohol screening and advice in an emergency department a nursing staff perspective. Accid Emerg Nurs 2007;15:3–9.
- [19] Glasgow RE, Bull SS, Piette JD, Steiner JF. Interactive behavior change technology. A partial solution to the competing demands of primary care. Am J Prev Med. 2004;27(2 Suppl 1):80–7.
- [20] Pinto BM, Friedman R, Marcus BH, Kelley H, Tennstedt S, Gillman MW. Effects of a computer-based, telephone-counseling system of physical activity. Am J Prev Med 2002;23:113–20.
- [21] Haerens L, Deforche B, Vandelanotte C, Maes L, De Bourdeaudhuij I. Acceptability, feasibility and effectiveness of a computer-tailored physical activity intervention in adolescents. Patient Educ Couns 2007;66:303–10.
- [22] Tate DF, Wing RR, Winett RA. Using Internet technology to deliver a behavioral weight loss program. JAMA 2001;285:1172–7.
- [23] Robinson R, West R. A comparison of computer and questionnaire methods of history-taking in a genito-urinary clinic. Psychol Health1992;6:77–84.
- [24] Thomas R, Cahill J, Santilli L. Using an interactive computer game to increase skill and self-efficacy regarding safer sex negotiation: field test results. Health Educ Behav 1997;24:71–86.
- [25] Locke SE, Kowaloff HB, Hoff RG, Safran C, Popovsky MA, Cotton DJ, Finkelstein DM, Page PL, Slack WV. Computer-based interview for screening blood donors for risk of HIV transmission. JAMA 1992;268:1301–5.

- [26] Tourangeau R, Smith TW. Asking sensitive questions. The impact of data collection mode, question format, and question context. Public Opin Q 1996;60:275–304.
- [27] MacMillan HL. Computer survey technology: a window on sensitive issues. Can Med Assoc J 1999;161:1142.
- [28] Noell J, Glasgow RE. Interactive technology applications for behavioral counseling.

 Issues and opportunities for health care settings. Am J Prev Med 1999;17:269–74.
- [29] Karlsson A, Johansson K, Nordqvist C, Bendtsen P. Feasibility of a computerized alcohol screening and personalized written advice in the ED: opportunities and obstacles. Accid Emerg Nurs 2005;13:44–53.
- [30] Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C, Buchner D, Ettinger W, Heath GW, King AC, Kriska A, Leon AS, Marcus BH, Morris J, Paffenbarger RS, Patrick K, Pollock ML, Rippe JM, Sallis J, Wilmore JH. Physical activity and public health: a recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. JAMA 1995;273:402–7.
- [31] Prochaska JO, DiClemente CC. Stages and processes of self change in smoking: toward an integrative model of change. J Consult Clin Psychol 1983;51:390–5.
- [32] Woolf SH, Krist AH, Johnson RE, Wilson DB, Rothemich SF, Norman GJ, Devers KJ.

 A practice-sponsored web site to help patients pursue healthy behaviors: an ACORN study. Ann Fam Med 2006;4:148–52.
- [33] Greenhalgh T, Robert G, Bate P, Macfarlane F, Kyriakidou O. Diffusion of innovations in health service organisations. A systematic literature review. Oxford: BMJ Books, Blackwell Publishing; 2005.

- [34] Beich A, Gannik D, Saelan H, Thorsen T. Screening and brief intervention targeting risky drinkers in Danish general practice a pragmatic controlled trial. Alcohol Alcohol 2007;42:593–603.
- [35] Bergman H, Källmén H. Alcohol use among Swedes and a psychometric evaluation of the alcohol use disorders identification test. Alcohol Alcohol 2002;37:245–51.
- [36] Källmén H, Wennberg P, Berman AH, Bergman H. Alcohol habits in Sweden during 1997–2005 measured with the AUDIT. Nord J Psychiatry 2007;61:466–70.
- [37] SNIPH Folkhälsopolitisk rapport 2005. Stockholm: Swedish National Institute of Public Health; 2005 (in Swedish, summary in English).
- [38] Aittasalo M, Miilunpalo S, Ståhl T, Kukkonen-Harjula K. From innovation to practice: initiation, implementation and evaluation of a physician-based physical activity promotion programme in Finland. Health Promot Int 2007;22:19–27.
- [39] Dugdill L, Graham RC, McNair F. Exercise referral: the public health panacea for physical activity promotion? A critical perspective of exercise referral schemes; their development and evaluation. Ergonomics 2005;48:1390–410.
- [40] Mikkelsen J, Holm HA. Contextual learning to improve health care and patient safety.

 Educ Health (Abingdon) 2007;20:124.