

## Alcohol and non-communicable diseases (NCDs): time for a serious international public health effort

*Together with smoking, diet and physical inactivity, consumption of alcohol is among the four most important risk factors for non-communicable disease (NCD). Alcohol consumption, especially heavy consumption, impacts on cancer, liver cirrhosis and stroke. To reduce the burden of NCD, effective alcohol policies should be implemented locally, nationally and internationally.*

Each year, in connection with the opening of the General Assembly sessions, the United Nations holds a special session on an agreed-upon topic. In 2011, the topic is non-communicable diseases (NCDs), arguing for their inclusion in the Millennium Development Goals, with the primary goal of emphasizing the importance of addressing such diseases in order to reduce the global burden of illness, not only in rich countries, but also in poor countries as a consequence of major epidemiological transitions [1–3].

While NCDs as a category covers a broad range of illnesses, primary emphasis has been put on cancer, heart disease, chronic respiratory diseases and diabetes [4]. Despite the fact that each disease has its own specific aetiology, there are risk factors which reach across many of the major NCDs including, in particular, several behavioural risk factors: unhealthy diet, lack of exercise, tobacco smoking and alcohol use.

One approach to limiting the play of such behavioural risk factors is based on a perspective of individual choice and responsibility. The premise is that educated consumers will act in their own long-term interests, even in the face of heavy marketing and promotion; controlling behaviour in such an environment may even be regarded as a test of virtue [5]. A perspective that educating the consumer is all that is needed has been the approach in recent times globally, not only in market economies. Many low- and middle-income countries have also adopted this approach, especially the emerging economies in Asia, with high economic growth rates. The experience of the second half of the 20th century was that such *laissez-faire* approaches are not a successful strategy for limiting NCDs.

A second approach has been to proscribe particular forms of consumption altogether. While this has obvious difficulties in the case of diet, for tobacco and alcohol there have been prohibitions on use in various forms and times. Again, the experience has been cautionary; such

prohibitions tend to succeed only when backed up strongly by religion and culture.

A third approach is through regulating the market to channel and influence consumer behaviour towards restrained and less harmful use [6]. Addressing such ‘upstream’ factors can be quite effective, as has been shown for alcohol and tobacco [7,8], but this strategy of balancing conflicting interests is inherently unstable. Because it involves tolerating a certain level of NCDs and other harm from the behaviour, it involves compromises and is easily seen as unprincipled. Regulating the actions of private interests in the market became more difficult in the late 20th century, in an era of globalization, free trade agreements and free market ideology [7, p. 71–101, 9]. Given the political influence of market interests, substantial and sustained counter-pressure from public health interests is required to prevent regulatory capture.

A regulatory approach thus requires substantial and sustained efforts in the public health interest. It will not be enough for the United Nations to pass a resolution at the Special Session and leave it at that. This is an issue for all the behavioural risk factors for NCDs, but it is especially an issue for alcohol. Alcohol is now included regularly in NCD discussions, such as the preparatory meeting for the September session in Moscow in April 2011 [10], but it regularly receives the least attention of the major risk factors. There are several reasons for this. For instance, the health-protective effects of alcohol in ischaemic disease for the middle-aged and older people [11] has confused and inhibited efforts to curtail harmful use of alcohol, even though the net effect of alcohol on cardiovascular disease—setting all else aside—is negative. Furthermore, moving on alcohol is more difficult because it is so much more part of the daily life of decision-makers: the affluent and powerful regularly use alcohol, whereas smoking tobacco or being overweight are increasingly characteristics of the poor and socially excluded. Finally, alcohol industry interests operate effectively in political spheres to minimize the efforts of public health proponents to address the impact of alcohol use on NCDs, among other harms. This occurs both at international levels, with increased global concentration in the spirits and beer industries [12], and nationally and locally, with widespread nets of producers and distributors—whether cider makers and whisky distillers in the United Kingdom, vintners in Australia and southern Europe, fruit wine producers in Finland or brewers in Belgium, India and China.

Against these forces the public health effort on alcohol is weak, particularly at the international level. To counter the pressures from globalized industries and from free-trade treaties and settlements, there is no Framework Convention on Alcohol Control [13] with a secretariat to increase its effectiveness, as there is for tobacco. As a first step forward, there is a new World Health Organization (WHO) Global Strategy on Alcohol [14] which sets out an initial framework for action on an international basis, but the resources available for implementing the strategy are puny. Whether as a reflection of behind-the-scenes pressure from the alcohol industry to starve the effort, of the ambivalence of political classes about alcohol, or of a withering of national commitments to international aid, the WHO alcohol programme is forced to operate on a shoestring budget, with minimal staff and programme resources.

Meanwhile, recognition of the necessity of addressing harmful use of alcohol in order to attain public health objectives increases. Alcohol plays an important causal role in the aetiology of NCDs. While this is particularly the case in countries in the former Soviet Union [3], the problems extend across the globe. Alcohol is the third leading risk factor in the global burden of disease for death and disability in general [15], and this is calculated primarily in terms of the adverse effects of drinking on the drinker. There are also large adverse effects of drinking on others [16], not counted in the health statistics. The case for increased priority being given to act on alcohol to address NCDs and other public health concerns is now very strong. In this context, the relative lack of action on alcohol is increasingly indefensible. The September meeting at the United Nations is an occasion for remedying this, and for taking concrete steps to increase the resources at the international level devoted to alcohol policy issues.

**Keywords** Alcohol, alcohol policy, international control, non-communicable disease (NCD), risk factor.

**Declarations of interest:**

None.

- 1 ROBIN ROOM,<sup>1</sup> JÜRGEN REHM<sup>2</sup> & CHARLES PARRY<sup>3</sup>  
 2 AER Centre for Alcohol Policy Research, Turning Point  
 3 Alcohol & Drug Centre, Fitzroy, Vic., Australia; School of  
 4 Population Health, University of Melbourne, Australia and  
 5 Centre for Social Research on Alcohol and Drugs, Stockholm  
 6 University, Sweden<sup>1</sup>, Centre for Addiction and Mental  
 7 Health, Toronto, Canada; Dalla Lana School of Public  
 8 Health and Department of Psychiatry, University of  
 9 Toronto, Canada and Institut für Klinische Psychologie und  
 10 Psychotherapie, Technische Universität Dresden, Germany<sup>2</sup>,  
 11 Alcohol and Drug Abuse Research Unit, Medical Research

Council, South Africa; Department of Psychiatry,  
 Stellenbosch University, South Africa and Global Alcohol  
 Policy Alliance<sup>3</sup>. E-mail: robinr@turningpoint.org.au

**References**

1. Omran A. R. The epidemiologic transition: a theory of the epidemiology of population change. *Milbank Mem Fund Q* 1971; 49: 509–38.
2. Room R., Rehm J. Alcohol and non-communicable diseases—cancer, heart disease and more. *Addiction* 2011; 106: 1–2.
3. Parry C., Patra J., Rehm J. Alcohol consumption and non-communicable diseases: epidemiology and policy implications. *Addiction*.
4. Alwan A. editor. *Global Status Report on Noncommunicable Diseases 2010*. Geneva: World Health Organization; 2011.
5. Room R. Addiction and personal responsibility as solutions to the contradictions of neoliberal consumerism. *Crit Public Health*, 2011; early view publication. DOI: 10.1080/09581596.2010.529424.
6. Thaler R. H., Sunstein C. R. *Nudge: Improving Decisions about Health, Wealth and Happiness*. New Haven/London: Yale University Press; 2008.
7. Babor T., Caetano R., Casswell S., Edwards G., Giesbrecht N., Graham, K. et al. *Alcohol: No Ordinary Commodity—Research and Public Policy*, 2nd edn. Oxford: Oxford University Press; 2010.
8. Chaloupka F., Strait K., Leon M. E. Effectiveness of tax and price policies in tobacco control. *Tob Control* 2011; 20: 235–8.
9. McGrady B. *Trade and Public Health: The WTO, Tobacco, Alcohol, and Diet*. Cambridge: Cambridge University Press; 2011.
10. First Global Ministerial Conference on Healthy Lifestyles and Noncommunicable Disease Control, Moscow, 28–29 April 2011. *Moscow Declaration*. Moscow: World Health Organization and Ministry of Public Health and Social Development, Russian Federation. Available at: [http://www.who.int/nmh/events/un\\_ncd\\_summit2011/en/index.html](http://www.who.int/nmh/events/un_ncd_summit2011/en/index.html) (accessed 11 June 2011).
11. Rehm J., Roerecke M. Alcohol, the heart and the cardiovascular system—what do we know and where should we go? *Drug Alcohol Rev* in press.
12. Jernigan D. The extent of global alcohol marketing and its impact on youth. *Contemp Drug Problems* 2010; 37: 57–89.
13. Room R., Schmidt L., Rehm J., Mäkelä P. International regulation of alcohol. *BMJ* 2008; 337: a2364.
14. World Health Organization (WHO). *Global Strategy to Reduce the Harmful Use of Alcohol*. Geneva: World Health Organization; (2010). Available at: [http://www.who.int/substance\\_abuse/activities/globalstrategy/en/index.html](http://www.who.int/substance_abuse/activities/globalstrategy/en/index.html) (accessed 10 June 2011).
15. World Health Organization (WHO). *Global Health Risks: Mortality and Burden of Disease Attributable to Selected Major Risks*. Geneva: World Health Organization; 2009. Available at: [http://www.who.int/healthinfo/global\\_burden\\_disease/global\\_health\\_risks/en/index.html](http://www.who.int/healthinfo/global_burden_disease/global_health_risks/en/index.html) (accessed 10 June 2011).
16. Laslett A.-M., Room R., Ferris J., Wilkinson C., Livingston M., Mugavin J. Surveying the range and magnitude of alcohol's harm to others in Australia. *Addiction*, in press.

<b>Toppan Best-set Premedia Limited</b>	
Journal Code: ADD	Proofreader: Jason
Article No: 3549	Delivery date: 11 July 2011
Page Extent: 2	

## AUTHOR QUERY FORM

Dear Author,

During the preparation of your manuscript for publication, the questions listed below have arisen. Please attend to these matters and return this form with your proof.

Many thanks for your assistance.

Query References	Query	Remark
q1	AUTHOR: Please advise if affiliations are correct	
q2	AUTHOR: please provide location of Global Alcohol Policy Alliance.	
q3	AUTHOR: AUTHOR: Authors are required to archive any web references before citing them using WebCite® technology ( <a href="http://www.webcitation.org">http://www.webcitation.org</a> ). This is an entirely free service that ensures that cited web material will remain available to readers in the future. Please provide the archived web reference, alongside the original URL, when returning proofs, e.g. (Archived at <a href="http://www.webcitation.org/5Kt3Pxf1">http://www.webcitation.org/5Kt3Pxf1</a> ).	
q4	WILEY-BLACKWELL: please complete when known.	
q5	AUTHOR: please complete reference if known.	
q6	AUTHOR: Reference 9 has not been included in the Reference List, please supply full publication details.	
q7	AUTHOR: please complete reference if known.	
q8	WILEY-BLACKWELL: please complete when known.	

# MARKED PROOF

## Please correct and return this set

Please use the proof correction marks shown below for all alterations and corrections. If you wish to return your proof by fax you should ensure that all amendments are written clearly in dark ink and are made well within the page margins.

<i>Instruction to printer</i>	<i>Textual mark</i>	<i>Marginal mark</i>
Leave unchanged	... under matter to remain	Ⓟ
Insert in text the matter indicated in the margin	∧	New matter followed by ∧ or ∧ <sup>Ⓢ</sup>
Delete	/ through single character, rule or underline or ┌───┐ through all characters to be deleted	Ⓞ or Ⓞ <sup>Ⓢ</sup>
Substitute character or substitute part of one or more word(s)	/ through letter or ┌───┐ through characters	new character / or new characters /
Change to italics	— under matter to be changed	↵
Change to capitals	≡ under matter to be changed	≡
Change to small capitals	≡ under matter to be changed	≡
Change to bold type	~ under matter to be changed	~
Change to bold italic	≈ under matter to be changed	≈
Change to lower case	Encircle matter to be changed	≡
Change italic to upright type	(As above)	⊕
Change bold to non-bold type	(As above)	⊖
Insert 'superior' character	/ through character or ∧ where required	Υ or Υ under character e.g. Υ or Υ
Insert 'inferior' character	(As above)	∧ over character e.g. ∧
Insert full stop	(As above)	⊙
Insert comma	(As above)	,
Insert single quotation marks	(As above)	ʹ or ʸ and/or ʹ or ʸ
Insert double quotation marks	(As above)	“ or ” and/or ” or ”
Insert hyphen	(As above)	⊥
Start new paragraph	┌	┌
No new paragraph	┐	┐
Transpose	└┐	└┐
Close up	linking ○ characters	Ⓞ
Insert or substitute space between characters or words	/ through character or ∧ where required	Υ
Reduce space between characters or words		↑